



0240

#5  
0300  
08264  
0220

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE  
Re the application of:

LUTTICKEN ET AL.

Serial Number: 09/084,837

Group Art Unit: to be assigned

Filed: May 26, 1998

Examiner: to be assigned

For: RECOMBINANT BIRNAVIRUS VACCINE

CLAIM TO PRIORITY UNDER 35 USC 119

Assistant Commissioner of Patents  
Washington, D.C. 20231

June 9, 1998

Sir:

The benefit of the filing date of the following prior foreign application is hereby requested for the above-identified application, and the priority provided in 35 USC 119 is hereby claimed:

**European Patent Application No. 972015994 filed May 26, 1997**

In support of this claim, the requisite certified copy of said original foreign application is filed herewith.

It is requested that the file of this application be marked to indicate that the Applicant has complied with the requirements of 35 USC 119 and that the Patent and Trademark Office kindly acknowledge receipt of this document.

In the event any fees are required with this paper, please charge our Deposit Account No. 02-2334.

Respectfully submitted,

*Mary E. Gormley*  
Mary E. Gormley  
Agent for Applicants  
Registration No. 34,409


Attorney Docket No. I/97269 US

AKZO NOBEL N.V.  
1300 Piccard Drive, Suite 206  
Rockville, Maryland 20850-4373  
Tel: (301) 948-7400 Fax: (301) 948-9751  
MEG/ms Enclosures  
68lutckn.pri

98 JUL 22 AM 9:35  
GROUP 180



Europäisches  
Patentamt



European  
Patent Office

Office européen  
des brevets

Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-  
gen stimmen mit der  
ursprünglich eingereichten  
Fassung der auf dem näch-  
sten Blatt bezeichneten  
europäischen Patentanmel-  
dung überein.

The attached documents  
are exact copies of the  
European patent application  
described on the following  
page, as originally filed.

Les documents fixés à  
cette attestation sont  
conformes à la version  
initialement déposée de  
la demande de brevet  
européen spécifiée à la  
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

97201599.4

CERTIFIED COPY OF  
PRIORITY DOCUMENT

RIBBON CUT BY  
CERTIFICATION BRANCH

RECEIVED  
98 JUL 22 AM 9:35  
GROUP 180

Der Präsident des Europäischen Patentamts:  
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets  
p.o.

M. B. RIJLING

DEN HAAG  
THE HAGUE 05/98  
LA HAYE, LE

EPA/EPO/OEB Form - 02.91



Europäisches  
Patentamt

European  
Patent Office

Office européen  
des brevets

**Blatt 2 der Bescheinigung**  
**Sheet 2 of the certificate**  
**Page 2 de l'attestation**

Anmeldung Nr.:  
Application no.:  
Demande n°: 97201599.4

Anmeldetag:  
Date of filing: 26/05/97  
Date de dépôt:

Anmelder:  
Applicant(s):  
Demandeur(s):  
Akzo Nobel N.V.  
6824 BM Arnhem  
NETHERLANDS

Bezeichnung der Erfindung:  
Title of the invention:  
Titre de l'invention:  
Recombinant birnavirus vaccine

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s) revendiquée(s)

Staat:  
State:  
Pays:

Tag:  
Date:  
Date:

Aktenzeichen:  
File no.  
Numéro de dépôt:

Internationale Patentklassifikation:  
International Patent classification:  
Classification internationale des brevets:  
C12N15/40, C12N7/01, A61K39/12, G01N33/569

Am Anmeldetag benannte Vertragsstaaten:  
Contracting states designated at date of filing: AT/BE/CH/DE/DK/ES/FI/FR/GB/GR/IE/IT/LI/LU/MC/NL/PT/SE  
Etats contractants désignés lors du dépôt:

Bemerkungen:  
Remarks:  
Remarques:

## Recombinant birnavirus vaccine

The present invention is concerned with a birnavirus mutant, a vaccine comprising this mutant, a method for determining birnavirus infection in an animal, as well as with a test kit for carrying out this method.

Infectious bursal disease virus (IBDV) and Infectious pancreatic necrosis virus (IPNV) are members of the Birnaviridae family. Viruses in this family have a very similar genomic organisation and a similar replication cycle. The genomes of these viruses consist of 2 segments (A and B) of double-stranded (ds) RNA. The larger segment A encodes a polyprotein which is cleaved by autoproteolysis to form mature viral proteins VP2, VP3 and VP4 (Hudson, P.J. et al, Nucleic Acids Res., 14, 5001-50012, 1986; Dobos P., Annual review of fish diseases 5, 25-54, 1995). VP2 and VP3 are the major structural proteins of the virion. VP2 is the major host-protective immunogen of birnaviruses, and contains the antigenic regions responsible for the induction of neutralising antibodies. The VP4 protein appears to be a virus-coded protease that is involved in the processing of a precursor polyprotein of the VP2, VP3 and VP4 proteins. The larger segment A possesses also a second open reading frame (ORF), preceding and partially overlapping the polyprotein gene. This second open reading frame encodes a protein VP5 of unknown function that is present in IBDV infected cells (Mundt, E. et al., J. Gen. Virol., 76, 437-443, 1995).

The smaller segment B encodes VP1, a 90 kDa multifunctional protein with polymerase and capping enzyme activities (Spies, U. et al., Virus Res., 8, 127-140, 1987 and Spies, U. et al., J. Gen. Virol., 71, 977-981, 1990; Duncan R. et al., Virology 181, 541-552, 1991).

For IBDV, two serotypes exist, serotype 1 and 2. The 2 serotypes may be differentiated by virus neutralisation (VN) tests. Furthermore, subtypes of serotype 1 have been isolated. These so-called "variant" viruses of serotype 1 can be identified by cross-neutralisation tests (Diseases of Poultry, 9th edition, 1991, Wolfe Publishing Ltd, ISBN 0 7234 1706 7, Chapter 28, P.D. Lukert and Y.M. Saif, 648-663), a panel of monoclonal antibodies (Snyder, D.B. et al., Arch. Virol., 127, 89-101. 1992.) or RT-PCR (Jackwood, D.J., Proceedings of the International symposium on infectious bursal disease and chicken infectious anaemia, Rauschholzhausen, Germany, 155-161, 1994). Some of these subtypes of serotype 1 of IBDV have been described in literature for example: Classical, Variant-E, GLS, RS593 and DS326 strains (Van Loon, et al. Proceedings of

the International symposium on infectious bursal disease and chicken infectious anaemia, Rauschholzhausen, Germany, 179-187, 1994).

5 Infectious Bursal disease (IBD), also called Gumboro disease, is an acute, highly-  
contagious viral infection in chickens that has lymphoid tissue as its primary target with a  
selective tropism for cells of the bursa of Fabricius. The morbidity rate in susceptible flocks is  
high, with rapid weight loss and moderate mortality rates. Chicks that recover from the disease  
may have immune deficiencies because of the destruction of the bursa of Fabricius which is  
essential to the defence mechanism of the chicken. The IBD-virus causes severe  
10 immunosuppression in chickens younger than 3 weeks of age and induces bursal lesions in  
chicks up to 3 months old.

For many years the disease could be prevented by inducing high levels of antibodies in  
breeder flocks by the application of an inactivated vaccine, to chickens that had been primed with  
15 attenuated live IBDV vaccine. This has kept economic losses caused by IBD to a minimum.  
Maternal antibodies in chickens derived from vaccinated breeders prevents early infection with  
IBDV and diminishes problems associated with immunosuppression. In addition, attenuated live  
vaccines have also been used successfully in commercial chicken flocks after maternal antibodies  
had declined.

20 Recently, very virulent strains of IBDV have caused outbreaks of disease with high  
mortality in Europe. The current vaccination programs failed to protect chicks sufficiently.  
Vaccination failures were mainly due to the inability of live vaccines to infect the birds before  
challenge with virulent field virus.

Eradication of the disease by other preventative measures than vaccination has not been  
25 feasible, because the virus is widely spread and because with currently administered live  
attenuated or inactivated IBDV vaccines it is not possible to determine whether a specific animal  
is infected with an IBDV field virus or whether the animal was vaccinated with an IBDV  
vaccine. In order to be able to start an eradication control programme for IBDV it is highly  
desirable that the possibility exists to discriminate between animals vaccinated with an IBDV  
30 vaccine and those infected with a field virus so as to be able to take appropriate measures, i.e.  
remove infected flocks, to reduce spreading of the virulent field virus. The introduction of, for  
example, a serologically identifiable marker can be achieved by introducing a mutation in genes

encoding non-essential (glyco)proteins of the IBDV which still give rise to the production of antibodies in an infected host animal. A marker vaccine for Aujeszky's disease and companion diagnostic tests have proven their practical value in the control of this disease. Whereas such control programs for other viral infectious diseases in animals are under development, until the present invention a vaccine based on an IBDV vaccine strain which would fit in IBDV control programs has not been described yet. The main reason for this is that the prerequisites for the development for such an IBDV marker vaccine were not met. No permissive position or region in the genomic IBDV sequence, i.e. a position or region which can be used for the incorporation of the mutation without disrupting essential functions of IBDV, such as those necessary for infection and replication, have been identified yet. Moreover, such a non-essential region in the IBDV genome should encode a (glyco)protein which elicits a major serological response in an animal infected with wild-type IBDV, and such a region was not identified before.

The present inventors have unexpectedly found a non-essential gene within segment A of a birnavirus genome which can be mutated such that the resulting birnavirus mutant does not produce the native expression product of that gene. Moreover, it has been found that this birnavirus mutant can be used as a marker vaccine virus which allows to make a serological distinction between animals infected with wild-type birnavirus and animals immunised with a vaccine based on this birnavirus mutant.

The present invention provides a birnavirus mutant which is not able to produce a native VP5 protein as a result of a mutation in the VP5 gene of the birnavirus genome.

Preferably, the birnavirus mutant is an IBDV mutant or an IPNV mutant, the IBDV mutant being most preferred.

The inventors have found that an IBDV mutant which is not able to produce the native VP5 protein is still able to infect cells and to replicate in these cells in vitro. It is demonstrated that the IBDV mutant according to the invention is replication competent in cell culture (Example 2). The VP5<sup>-</sup> IBDV exhibits a delay in replication in chicken embryo cells as compared to the VP5<sup>+</sup> parental virus, however, final yields of the virus are similar, i.e. about 10<sup>7.5</sup> TCID<sub>50</sub>/ml (Example 1). Moreover, it is demonstrated that the IBDV mutant is also able to infect poultry and to replicate in the infected host animals in vivo, i.e. evidence is provided that the gene encoding the VP5 protein is a non-essential gene. Example 3 shows that the VP5<sup>-</sup> IBDV can be re-isolated

from organs of animals infected with the IBDV mutant and that the IBDV mutant induces a protective immune response in the infected animals.

Moreover, it has been established herein that part of the normal anti-IBDV immune response in poultry is directed to the VP5 region. This is rather surprising as the VP5 protein is considered to represent a non-structural viral protein (Mundt et al., J. Gen. Virol. 76, 437-443, 1995) and the immune response in an animal against a viral pathogen is usually elicited against the structural (glyco)proteins of the virus. These findings make the IBDV mutant and other birnavirus mutants according to the present invention a suitable vaccine candidate for a marker vaccine. Such a marker vaccine provides the possibility to determine whether animals are infected with a wild-type birnavirus, e.g. IBDV, or with a vaccine virus.

Additionally, it has been found that the VP5 protein is involved in the expression of virulence of the birnaviruses, in particular of IBDV, and that the inability of the virus mutants to produce the native VP5 protein leads to an attenuation of the virus.

With the term "which is not able to produce a native VP5 protein" is meant that the birnavirus mutant produces a polypeptide that can be distinguished by serological tests from the native VP5 protein, or does not produce a VP5 protein at all. For example, in the former case, the birnavirus mutant produces only a fragment of the native birnavirus VP5 protein which lacks one or more immunogenic epitopes.

Preferably, the birnavirus mutant according to the invention produces no VP5 protein upon infection of a host cell.

As described above, the genomic organisation of the birnaviruses is well established: the IBDV and IPNV genome comprises a large segment A and a smaller segment B. The segment A of IBDV comprises a large open reading frame (ORF) encoding a polyprotein of about 110 kDa (VP2-VP4-VP3). The gene encoding the VP5 protein is identified in the prior art, and defined herein, as the small ORF on segment A of the birnavirus genome which precedes and partially overlaps the polyprotein encoding ORF (Bayliss et al., J. Gen. Virol. 71, 1303-1312, 1990; Spies et al., J. Gen. Virol. 71, 977-981, 1990; Havarstein L.S. et al., J. Gen. Virology 71, 299-308; 1990; Dobos et al., 1995, supra; Figures 1-3 herein and SEQ ID No.'s 1-7). The mutation introduced in the VP5 gene is such that it does not prevent the expression of the polyprotein.

SEQ ID No. 1 comprises the full length cDNA nucleotide sequence of segment B of IBDV strain P2, as well as the amino acid sequence of the VP1 protein encoded by segment B (see also SEQ ID. No. 2). SEQ ID No. 3 and 5 depict the full length cDNA sequence of segment A of

IBDV strain D78 and the coding region of the VP5 protein and the polyprotein, respectively. SEQ ID 3 and 4 also show the amino acid sequence of the D78 VP5 protein. SEQ ID No. 5 and 6 show the amino acid sequence of the polyprotein VP2-VP4-VP3 of D78. SEQ ID No. 7 shows the 5'-end of segment A of strain D78, including the mutations introduced in the VP5 coding region. SEQ ID No. 8 shows the nucleotide sequence of segment B of strain D78 and the amino acid sequence of the D78 VP1 protein. The genomic organisation of both segments is also shown in Figure 1.

The ORF coding for VP5 is conserved in all hitherto published segment A sequences. The IBDV ORF encodes 145 amino acids resulting in a calculated molecular mass of 16.5 kDa. The nucleotide sequence of the ORF encoding the VP5 protein of IBDV strain D78 used herein is shown in SEQ ID No. 3 and 4. Natural variations may exist between individual IBDV isolates. These natural variations result from small differences in the genomes of these viruses. The nucleotide sequence of the segment A, including the nucleotide sequence of the VP5 gene for many IBDV isolates have been described in the prior art (Vakharia et al., *Avian Diseases* 36, 736-742, 1992; Bayliss et al., *J. Gen. Virol.* 71, 1303-1314, 1990; Hudson et al., *Nuc. Acid Res.* 14, 5001-5012, 1986; Schnitzler et al., *J. Gen. Virol.* 47, 1563-1571, 1993; Kibenge et al., *J. Gen. Virol.* 71, 569-577, 1990 and *Virology* 184, 437-440, 1991; Mundt et al., *Virology* 209, 10-18, 1995; Lana et al., *Virus Genes* 6, 247-259, 1992; Vakharia et al., *Virus Res.* 31, 265-273, 1994; Brown et al., *Virus Res.* 40, 1-15, 1996). The amino acid sequence of the VP5 protein from serotype I IBDV strains display a homology of at least 95% with the VP5 amino acid sequence shown in SEQ ID No. 3 and 4, whereas the homology between serotype II VP5 sequence and the amino acid sequence shown in SEQ ID No. 3 and 4 is at least 75%. Therefore, a preferred IBDV mutant according to the present invention is an IBDV mutant wherein the mutation is introduced in the VP5 gene having a homology of at least 75%, in particular at least 95% on the amino acid sequence level with the VP5 amino acid sequence shown herein.

Preferably an IBDV mutant according to the present invention is derived from any of the classical or variant (e.g. variant E or GLS) IBDV vaccine strains, such as those currently used in the field. Such suitable IBDV strains include the IBDV vaccine strains present in the commercially available vaccines: D78, PBG 98, LZ 228E, 89-03 (Intervet International B.V.), Bursine 2 (Fort Dodge Animal Health) and S 706 (Rhône Mérieux).



A particular preferred IBDV mutant according to the invention is derived from the D78 strain comprising a VP5 gene encoding a protein having the amino acid sequence shown in SEQ ID No. 3 and 4.

Alternatively, the parent birnavirus strain for the virus mutant according to the invention is a virulent birnavirus field strain. It is found herein that the VP5 protein is a factor associated with virulence, and that the absence of the native VP5 protein in a birnavirus results in an attenuated form of the virus.

Preferably the invention provides a birnavirus mutant which is not able to produce a native VP5 protein as a result of a mutation in the part of the VP5 gene which does not overlap with the large ORF encoding the polyprotein.

In particular, the birnavirus mutant according to the invention comprises a mutation in the 5'-end of the VP5 gene spanning nucleotides 1-30, preferably 1-20, more preferably 1-10. Most preferred is an birnavirus mutant having a mutation in nucleotides 1-3 of the VP5 gene.

A mutation is understood to be a change of the genetic information in the VP5 gene with respect to the genetic information present in this region of the genome of naturally occurring birnavirus producing native VP5 protein. The mutation is, for example, a nucleic acid substitution, deletion, insertion or inversion, or a combination thereof.

In a preferred embodiment of the present invention a birnavirus mutant is provided wherein the mutation is a substitution of one or more nucleotides. In particular, a nucleic acid substitution is introduced in the start codon, as a result of which the new codon encodes an amino acid different from methionine or represents a stop codon, preferably the nucleic acid substitution comprises at least two of the nucleotides of the start codon.

A further birnavirus mutant according to the invention comprises a substitution of one or more nucleotides in a codon(s) different from the start codon resulting in one or more stop codons, preferably in the 5'-end of the VP5 gene as defined above, if desired in addition to a substitution in the start codon as described above. Preferably, the birnavirus mutant comprises a stop codon in this region of the VP5 gene in each of the three reading frames.

Such a preferred birnavirus mutant may be an IBDV mutant having a mutation in the start codon, the fourth and the sixth codon of the VP5 gene, preferably resulting in the mutated codons shown in SEQ ID No. 7 and Figure 3.

Alternatively, a birnavirus mutant is provided wherein the mutation is a deletion. In particular, the deletion comprises less than 20, less than 10 or less than 5 nucleotides. Preferably, the deletion comprises a total number of nucleotides not dividable by three, resulting in a shift of the reading frame.

5        Preferably the deletion comprises one or more nucleotides of the start codon of the VP5 gene.

10        In an alternative embodiment of the present invention a birnavirus mutant is provided wherein the mutation comprises the insertion of a heterologous nucleic acid sequence in the birnavirus genome. A heterologous nucleic acid sequence is a nucleic acid sequence normally not present at the specific insertion site of the particular virus species.

15        The heterologous nucleic sequence to be incorporated into the birnavirus genome is a nucleic acid fragment which either encodes a polypeptide or is a non-coding sequence. The nucleic acid fragment can be derived from any source, e.g. viral, eukaryotic, prokaryotic or synthetic, including oligonucleotides suitable for the interruption of the expression of the VP5 gene.

20        A suitable oligonucleotide for the interruption of the VP5 expression may comprise three translational stop codons in each of the possible reading frames in both directions, in addition to one or more appropriate restriction enzyme cleavage sites useful for the insertion of a second heterologous nucleic acid sequence. The length and nucleotide sequence of such a non-coding heterologous nucleic acid sequence is not critical, but preferably varies between 8-50 nucleotides.

25        In a further embodiment of the present invention a birnavirus mutant is provided which can be used not only for the preparation of a vaccine against infection by a specific birnavirus, but also against other poultry or fish infectious diseases. For example, a vector vaccine based on such an IBDV mutant offers the possibility to immunise against other avian pathogens by the expression of antigens of these avian pathogens within infected cells of the immunised host. Such an IBDV vector according to the present invention can be obtained by inserting a heterologous nucleic acid sequence encoding a polypeptide heterologous to the IBDV in the VP5 gene as defined herein.

30        The heterologous nucleic acid sequence may encode an antigen of an avian pathogen such as Newcastle disease virus, Infectious bronchitis virus, Marek's disease virus, avian encephalomyelitis virus, avian reovirus, avian influenza virus, chicken anaemia virus, *Salmonella* spp., *E.coli*, and *Eimeria* spp.

Furthermore, an IBDV mutant according to the invention comprises in addition to the mutation in the VP5 gene, a mutation in the VP2 gene, wherein this gene expresses a chimeric protein comprising neutralising epitopes of more than one antigenic type of IBDV (e.g. classic, Variant-E and/or GLS). Preferably, such a mutant comprises the relevant protective VP2 epitopes of a variant GLS strain and classic strain. In particular, the mutated VP2 gene is a GLS VP2 gene comprising a nucleic acid sequence fragment encoding the B69 epitope. The construction of such a mutated VP2 genes is described in Snyder et al., Avian Diseases 38, 701-707, 1994.

Furthermore, nucleic acid sequences encoding polypeptides for pharmaceutical or diagnostic applications, in particular immuno-modulators such as lymphokines, interferons or cytokines, may be incorporated into the VP5 gene. The heterologous nucleic acid sequence may also encode a screenable marker, such as *E. coli*  $\beta$ -galactosidase or *E. coli*  $\beta$ -glucuronidase.

The construction of birnavirus mutants, in particular of IBDV mutants according to the present invention can be achieved by means of the recently established infectious cRNA system for IBDV (Mundt and Vakharia, Proc. Natl. Acad. Sci. USA 93, 11131-11136, 1996). This reverse genetics system opens the possibility to introduce mutations in the RNA genome of the IBD virus, in particular in the VP5 gene. The most important step in this reverse genetics system is to provide full length cDNA clones of the segments A and B of IBD virus. cDNA constructs comprising the segment A or B, including the nucleotides of the 5'- and 3'- ends of both these segments can be generated according to the method described by Mundt and Vakharia (1996, supra). Additionally, these constructs comprise a RNA polymerase promoter operably linked to either of the segments. The promoter can be the promoter for the T7, SP6 or T3 polymerase, the T7 promoter being preferred. Mutations can be introduced into the VP5 gene by means of methods generally known in the art for this purpose. In particular, the mutation(s) are introduced by means of site directed mutagenesis.

For example, in a first step a cDNA fragment is provided comprising at least a substantial part of the VP5 gene. In the next step suitable primer pairs are designed and hybridised with the VP5 sequence containing fragment. The 5'-primer comprises in addition to sequences complementary to the VP5 sequence, nucleotides which harbour the desired mutation, e.g. a mutation which changes the ATG start codon to an AGG (arginine) codon. Moreover, the 5'-primer is provided with an upstream nucleotide sequence representing a suitable restriction enzyme cleavage site which allows the restoring of the complete 5'-end non-coding sequence. Subsequently, the new mutated fragment is amplified using PCR and the new fragment is

introduced in the starting sequence by replacing the native nucleic acid sequence using appropriate restriction enzymes. In the next step plus-sense transcripts of the segment A and B are generated in vitro with (T7) RNA polymerase, after which the synthetic transcripts are purified using conventional RNA purification techniques. The recombinant IBDV mutant according to the invention is obtained after transfection of suitable cells (e.g. VERO cells, QM-7 cells or CEC cells) with the synthetic RNA transcripts of both segments of the IBDV genome, if desired in the presence of transfection-enhancing compositions, such as Lipofectin. Finally the recombinant IBDV is harvested from the supernatant of the transformed cells.

Methods for introducing a mutation in the birnavirus genome are described herein, but are also generally used in the art (Mundt and Vakharia, 1996, supra; Current Protocols in Molecular Biology, eds.: F. M. Ausubel et al., Wiley N.Y., 1995 edition, pages 8.5.1.-8.5.9.)

Further to the unexpected finding by the present inventors that the VP5 ORF of IBDV is a non-essential region of the IBDV genome, it has also been found that an IBDV mutant according to the present invention is able to induce a protective immune response, i.e. animals immunised with a vaccine comprising the IBDV mutant are protected against virulent challenge. Moreover, it has been found that anti-sera of animals infected with naturally occurring IBDV comprise antibodies directed to the non-structural VP5 protein and that these antisera can be distinguished from anti-sera derived from animals infected with an IBDV mutant according to the present invention. In addition, it has been found that the IBDV mutant as described above is attenuated if compared with the parent IBD virus which is able to produce the native VP5 protein.

Therefore, another aspect of this invention is a vaccine for use in the protection of animals against birnavirus infection comprising the birnavirus mutant as characterised above, together with a pharmaceutical acceptable carrier or diluent. In particular, the vaccine according to the invention is a vaccine for use in the protection of poultry against infectious bursal disease comprising the IBDV mutant described above.

The birnavirus mutant according to the present invention can be incorporated into the vaccine as live or inactivated virus.

A vaccine according to the invention can be prepared by conventional methods such as for example commonly used for the commercially available live- and inactivated IBDV vaccines. Briefly, a susceptible substrate is inoculated with an IBDV mutant according to the invention and

propagated until the virus replicated to a desired infectious titre after which IBDV containing material is harvested.

Every substrate which is able to support the replication of IBD viruses can be used in the present invention, including primary (avian) cell cultures, such as chicken embryo fibroblast cells (CEF) or chicken kidney cells (CK), mammalian cell lines such as the VERO cell line or the BGM-70 cell line, or avian cell lines such as QT-35, QM-7 or LMH. Usually, after inoculation of the cells, the virus is propagated for 3-10 days, after which the cell culture supernatant is harvested, and if desired filtered or centrifuged in order to remove cell debris.

Alternatively, the IBDV mutant is propagated in embryonated chicken eggs. In particular, the substrate on which these IBD viruses are propagated are SPF embryonated eggs. Embryonated eggs can be inoculated with, for example 0.2 ml IBDV mutant containing suspension or homogenate comprising at least  $10^2$  TCID<sub>50</sub> per egg, and subsequently incubated at 37 °C. After about 2-5 days the IBD virus product can be harvested by collecting the embryo's and/or the membranes and/or the allantoic fluid followed by appropriate homogenising of this material. The homogenate can be centrifuged thereafter for 10 min at 2500 x g followed by filtering the supernatant through a filter (100 µm).

The vaccine according to the invention containing the live virus can be prepared and marketed in the form of a suspension or in a lyophilised form and additionally contains a pharmaceutically acceptable carrier or diluent customary used for such compositions. Carriers include stabilisers, preservatives and buffers. Suitable stabilisers are, for example SPGA, carbohydrates (such as sorbitol, mannitol, starch, sucrose, dextran, glutamate or glucose), proteins (such as dried milk serum, albumin or casein) or degradation products thereof. Suitable buffers are for example alkali metal phosphates. Suitable preservatives are thimerosal, merthiolate and gentamicin. Diluents include water, aqueous buffer (such as buffered saline), alcohols and polyols (such as glycerol).

If desired, the live vaccines according to the invention may contain an adjuvant. Examples of suitable compounds and compositions with adjuvant activity are the same as mentioned below.

Although administration by injection, e.g. intramuscular, subcutaneous of the live vaccine according to the present invention is possible, the vaccine is preferably administered by the inexpensive mass application techniques commonly used for IBDV vaccination. For IBDV vaccination these techniques include drinking water and spray vaccination.

Alternative methods for the administration of the live vaccine include eye drop and beak dipping administration.

In another aspect of the present invention a vaccine is provided comprising the birnavirus mutant in an inactivated form. The major advantage of an inactivated vaccine is the extremely high levels of protective antibodies of long duration that can be achieved.

The aim of inactivation of the viruses harvested after the propagation step is to eliminate reproduction of the viruses. In general, this can be achieved by chemical or physical means. Chemical inactivation can be effected by treating the viruses with, for example, enzymes, formaldehyde,  $\beta$ -propiolactone, ethylene-imine or a derivative thereof. If necessary, the inactivating compound is neutralised afterwards. Material inactivated with formaldehyde can, for example, be neutralised with thiosulphate. Physical inactivation can preferably be carried out by subjecting the viruses to energy-rich radiation, such as UV light or  $\gamma$ -rays. If desired, after treatment the pH can be adjusted to a value of about 7.

A vaccine containing the inactivated birnavirus mutant can, for example comprise one or more of the above-mentioned pharmaceutically acceptable carriers or diluents suited for this purpose.

Preferably, an inactivated vaccine according to the invention comprises one or more compounds with adjuvant activity. Suitable compounds or compositions for this purpose include aluminium hydroxide, -phosphate or -oxide, oil-in-water or water-in-oil emulsion based on, for example a mineral oil, such as Bayol F® or Marcol 52® or a vegetable oil such as vitamin E acetate, and saponins.

The vaccine according to the invention comprises an effective dosage of the birnavirus mutant as the active component, i.e. an amount of immunising birnavirus material that will induce immunity in the vaccinated birds against challenge by a virulent virus. Immunity is defined herein as the induction of a significant higher level of protection in a population of birds after vaccination compared to an unvaccinated group.

Typically, the live vaccine according to the invention can be administered in a dose of  $10^2$ - $10^9$  TCID<sub>50</sub> infectious dose<sub>50</sub> (TCID<sub>50</sub>) per animal, preferably in a dose ranging from  $10^{5.0}$ - $10^{7.0}$  TCID<sub>50</sub>, and an inactivated vaccines may contain the antigenic equivalent of  $10^5$ - $10^9$  TCID<sub>50</sub> per animal.

Inactivated vaccines are usually administered parenterally, e.g. intramuscularly or subcutaneously.

Although, the IBDV vaccine according to the present invention may be used effectively in chickens, also other poultry such as turkeys, guinea fowl and partridges may be successfully vaccinated with the vaccine. Chickens include broilers, reproduction stock and laying stock.

5 The age of the animals receiving a live or inactivated vaccine according to the invention is the same as that of the animals receiving the conventional live- or inactivated IBDV vaccines. For example, broilers (free of maternally derived antibodies-MDA) may be vaccinated at one-day-old, whereas broilers with high levels of MDA are preferably vaccinated at 2-3 weeks of age. Laying stock or reproduction stock with low levels of MDA may be vaccinated at 1-10 days of age followed by booster vaccinations with inactivated vaccine on 6-8 and 16-20 weeks of age.

10 The invention also includes combination vaccines comprising, in addition to the IBDV or IPNV mutant according to the invention, one or more immunogens derived from other pathogens infectious to poultry or fish, respectively.

Preferably, the combination vaccine additionally comprises one or more vaccine strains of infectious bronchitis virus (IBV), Newcastle disease virus (NDV), egg drop syndrome (EDS)  
15 virus, turkey rhinotracheitis virus (TRTV) or reovirus.

In addition to a marker vaccine for birnaviruses, the availability of an appropriate diagnostic test is an essential requirement for the application of a birnavirus eradication control programme. Such a diagnostic test is provided herein and comprises a method for determining  
20 IBDV infection in poultry and IPNV infection in fish, i.e. it provides a method for distinguishing an animal in the field vaccinated with a vaccine as described above, from an animal infected with a naturally-occurring IBDV or IPNV.

Therefore, the present invention provides a method for the detection of birnavirus infection, in particular for the detection of IBDV infection in an animal comprising the step of examining a  
25 sample of the animal for the presence of VP5 antibodies or antigens. The animal is an animal from the field and is in particular an avian species, preferably a chicken. The sample coming from the animal may be any sample in which IBDV antibodies or antigens are present, e.g. a blood, serum or tissue sample, the serum sample being preferred.

A preferred method for determining birnavirus infection in an animal is a method for the detection of antibodies against the VP5 protein, comprising the steps of:

- (i) incubating a sample suspected of containing anti-birnavirus antibodies, with VP5 antigen,
- (ii) allowing the formation of antibody-antigen complex , and
- 5 (ii) detecting the presence of the antibody-antigen complex.

The design of this immunoassay may vary. For example, the immunoassay may be based upon competition or direct reaction. Furthermore, protocols may use solid supports or may use cellular material. The detection of the antibody-antigen complex may involve the use of labelled antibodies; the labels may be, for example, enzymes, fluorescent-, chemiluminescent-, radio-  
10 active- or dye molecules.

Suitable methods for the detection of the VP5 antibodies in the sample include the enzyme-linked immunosorbent assay (ELISA), immunofluorescent test (IFT) and Western blot analysis.

In an exemplifying ELISA, the wells of a polystyrene micro-titration plate are coated with VP5 antigen. Next, the wells of the coated plates are filled with chicken serum and serial  
15 dilutions are made. After incubation, chicken anti-VP5 protein serum antibodies are determined by detecting antibody (monoclonal or polyclonal) with the same specificity as the coated one, but which is labelled (e.g. with biotin). The labelled antibody will occupy the free antigens that have not been occupied by anti-VP5 antibodies in the chicken serum. For example, horse radish peroxidase coupled to avidin may be added and the amount of peroxidase is measured by an  
20 enzymatic reaction. If no antibodies against VP5 are present in the chicken serum sample then a maximum absorption is obtained. If the serum contains many antibodies against VP5 then a low absorption is expected. Alternatively, after the incubation with chicken serum, the amount of antibodies present in the serum that bound to the VP5 antigen may be determined directly by using an anti-chicken conjugate followed by the enzymatic reaction.

25 In a sandwich ELISA the wells of a polystyrene micro-titration plate can be coated with a monoclonal antibody directed against the VP5 protein. Next, the wells of these coated plates are incubated with VP5 antigen. After the antigen is captured, the wells are filled with the chicken serum and serial dilutions are made. Subsequently, the protocol as described above may be followed. This test can also be carried out by using polyclonal serum against VP5 instead of the  
30 coated monoclonal antibodies.

In another diagnostic test (Western blot analysis), the VP5 antigen (containing) material is subjected to SDS-PAGE. Next, the separated proteins are electroblotted onto nitro-cellulose



membrane. Thereafter, the membranes can be cut into lanes and the lanes are incubated with the chicken serum. The presence of VP5 antibodies in the sample can be determined by examination whether antibodies bound to the VP5 antigen, for example by using an anti-chicken conjugate followed by an enzymatic reaction. If antibodies against VP5 are present then a band at about 17 kDa is identifiable.

The VP5 antigen may be any VP5 protein (fragment) comprising material which allows the formation of the VP5 antigen-VP5 antibody complex. Preferably, the VP5 antigen comprises the expression product of a conventional recombinant host cell or virus, e.g. such as E.coli expressed VP5 (Mundt et al., J. Gen. Virol. 76, 437-443, 1995) or baculovirus expressed protein (Vakharia et al., Vaccine 12, 452-456, 1994; Vakharia et al., J. Gen Virol. 74, 1201-1206, 1993). In a further embodiment of the present invention a diagnostic test kit is provided which is suitable for performing the diagnostic test according to the invention as described above.

In particular, a diagnostic test kit is provided which comprises in addition to the components usually present, the VP5 antigen (if desired coated onto a solid phase) as the immunological reagent. Other components usually present in such a test kit include, biotin or horseradish peroxidase conjugated antibodies, enzyme substrate, washing buffer etc.

To determine birnavirus VP5 antigen in a test sample from an animal in the field, VP5-specific antibodies are used as the immunological reagent, preferably fixed to a solid phase. The test sample is added, and after an incubation time allowing formation of the antibody-antigen complex, a second labelled antibody may be added to detect the complex.

## EXAMPLES

### Example 1.

#### Construction and analysis of recombinant VP5<sup>-</sup> IBD virus

#### **Construction of full length VP5<sup>-</sup> clone of IBDV segment A.**

To construct a VP5-negative IBDV, the *EcoRI* site immediately following the 3'-end of the full length cDNA of strain D78 segment A (pUC19FLAD78; Mundt and Vakharia, Proc. Natl. Acad. Sci. USA 93, 11131-11136, 1996) was deleted. An *EcoRI* - *KpnI* fragment containing the

T7 polymerase binding site followed by the complete segment A sequence was excised and inserted into *EcoRI* - *KpnI* cleaved vector pUC18 after inactivation of the unique *NdeI* within the vector sequence resulting in plasmid pAD78/EK. Thereafter, the genomic region encompassing the initiation codon for VP5 was amplified in two pieces using primers A1F5' and VP5MutR, and VP5MutF and A2R, respectively (see Table 1 for sequence and location of primers). PCR fragments were cloned separately and were subsequently fused via a unique *AflIII* site which had been created by mutations within respective primers (see Fig. 2). An *EcoRI* - *NdeI* fragment containing the T7 polymerase binding site, and the 5'-part of segment A including the introduced mutations was excised and used to substitute the wild-type *EcoRI* - *NdeI* fragment in pAD78/EK to yield plasmid pAD78/VP5<sup>-</sup>. Of the three mutations introduced one altered the initiation methionine codon for VP5 into an arginine codon (Fig. 2).

**Table 1:** Sequence of oligonucleotide primers used for generating mutant constructs.

<sup>a</sup> Nucleotide sequence	Orientation	Designation	Nucleotide no.
AGAGAATTCTAATACGACTCACTATAGGA TACGATCGGTCTGAC	+	A1F5'	1-18
TGGGCCTGTCAGTGTGTCACATGT	-	A2R	716 - 740
CATTGCTCTGCAGTGTGTAGTGAGC	-	A3R	338 - 362
CTACAACGCTATCCTTAAGGGTTAGTA GAG	+	VP5MutF	80 - 109
CTCTACTAACCCTTAAGGATAGCGTTGT AG	-	VP5MutR	80 - 109

- a) Underlined nucleotides denote virus specific nucleotides. T7 promotor sequences are marked in italics. Mutated nucleotides are bold and orientation of the primer is shown for sense (+) and antisense (-). Primer positions are given according to the published sequence of serotype I strain P2 (Mundt et al., *Virology* 209, 209-218, 1995).

**Virus recovery from cRNA.** For *in vitro* transcription of RNA plasmids pAD78/EK, pAD78/VP5<sup>-</sup> and pBP2 (Fig. 2) were linearized by cleavage with *BsrGI* and *PstI*, respectively. Treatment of linearized DNA, transcription and purification of RNA, and transfection were carried out as described by Mundt and Vakharia (1996, *supra*) with the exception that secondary

CEC were used for the transfection experiments. Three days after transfection a CPE was visible in CEC. Cells were freeze/thawed, centrifuged at 700 x g to eliminate cellular debris, and the resulting supernatants were filtrated through 0.45  $\mu$ m filters and stored at -20°C. For the transfection experiments full length cDNA clones of segment A of strain D78 capable of expressing (pAD78/EK) or unable to express VP5 (pAD78/VP5<sup>-</sup>) were transcribed into synthetic RNA and cotransfected with segment B full length cRNA into CEC. Resulting virus progeny IBDV/EK and IBDV/VP5<sup>-</sup> was further characterised.

**Analysis of transfection progeny by immunofluorescence and Radioimmunoprecipitation assay (RIPA).** VP5 was expressed in E.coli as described in Mundt et al. (J. Gen. Virol. 76, 437-443, 1995). Rabbit monospecific polyclonal anti serum and mouse monoclonal antibodies against VP5 were prepared according to standard protocols. Vero cells infected with IBDV/VP5<sup>-</sup>, IBDV/EK, and non-infected cells, respectively, were incubated with rabbit anti-IBDV serum, rabbit anti-VP5 serum and with anti-VP5 mAb DIE 7, and stained with fluoresceine-conjugated secondary antibodies. Both antisera and the monoclonal antibody recognised IBDV antigens in the cytoplasm of IBDV/EK infected cells. In contrast, whereas the anti-IBDV serum readily detected viral antigens in IBDV/VP5<sup>-</sup> infected cells, neither the monospecific anti VP5-serum nor the monoclonal anti-VP5 antibody exhibited specific reactivity. None of these immunological reagents reacted with non-infected controls.

To analyse viral proteins expressed during replication lysates of radioactively labelled CEC infected with IBDV/VP5<sup>-</sup> (Fig 4, lanes 1-3) and IBDV/EK (Fig. 4, lanes 4-6) were immunoprecipitated with rabbit anti-IBDV serum, rabbit anti-VP5 serum and mAb DIE 7. Non-infected CEC were used as control (Fig. 4, lanes 7-9). IBDV/EK (lane 4) as well as IBDV/VP5<sup>-</sup> (lane 1) infected CEC showed viral proteins VP2, VP3, and VP4 after precipitation with rabbit anti-IBDV serum. The rabbit anti-VP5 serum (lane 5) and mAb DIE 7 (lane 6) precipitated VP5 with a molecular mass of 21 kDa only from IBDV/EK infected cells. No specific reactivity was detectable in IBDV/VP5<sup>-</sup> infected CEC after precipitation with rabbit-anti VP5 (lane 2) as well as the VP5 specific mAb DIE 7 (lane 3). Non-infected CEC showed no specific reactivity (lanes 7-9).

**Replication of IBDV/VP5<sup>-</sup> in CEC.** To assay replication of IBDV/VP5<sup>-</sup> in more detail one step growth was analysed (Fig. 5). Confluent secondary CEC were infected with IBDV/EK and IBDV/VP5<sup>-</sup> with  $10^{7.2}$  TCID<sub>50</sub>, respectively. Immediately after overlaying the infected cells with 5 ml growth medium, supernatant from one infected CEC tissue plate of each virus was removed and stored at -20°C (0 h p.i.). Remaining tissue culture plates were further incubated and 4h, 8h, 16h, 24h, and 48h p.i. supernatants were removed and stored at -20°C. Supernatants were centrifuged and titrated according to standard methods. The TCID<sub>50</sub> at the different time points after infection showed that the VP5 expressing virus (IBDV/EK) replicated faster than the virus mutant lacking VP5 (IBDV/VP5<sup>-</sup>). 16 h after infection IBDV/EK showed a 100-fold higher than IBDV/VP5<sup>-</sup> (Fig. 5). However, at 48 h p.i. IBDV/VP5<sup>-</sup> reached a titre of  $10^{7.2}$  TCID<sub>50</sub>/ml which was similar to IBDV/EK ( $10^{7.45}$ /ml)

**Preparation of recombinant IBDV VP5<sup>-</sup>-2.** Plasmid pAD78/VP5<sup>-</sup>-2 was prepared by techniques similar to those described above. The nucleotide sequence of part of the mutated VP5 gene is shown in SEQ ID No. 7 and Figure 3. A restriction enzyme fragment harbouring the mutations was used to substitute the wild-type *Eco*RI - *Nde*I fragment in pAD78/EK. An outline of the protocol for the preparation of the recombinant plasmid is shown in Figure 3. The organisation of pBD78 is also depicted in Figure 3. The recombinant virus was prepared as described above, except for the fact that segment B of strain D78 (SEQ ID No. 8) was used and QM-7 cells were used for the transfection experiment.

## Example 2

### Identification of VP5 protein in different IBDV strains

Different strains of IBDV were investigated for the expression of the VP5-gene. This was done by making use of the immuno-fluorescence technique (IFT). Chicken embryo fibroblasts grown in microtiterplates were infected with different IBDV strains. Three to 5 days after incubation at 37°C cells were fixed with 70% ethanol, then treated with polyclonal rabbit anti IBDV serum (R1928), polyclonal rabbit anti VP5 serum (RαVP5) or monoclonal antibody directed against VP5 (DIE7), respectively. Binding of the poly- or monoclonal antibodies to the

different IBDV strains was visualised by making use of a fluorescence labelled conjugate (goat-anti-rabbit or goat-anti-mouse). The results are shown in Table 2:

5 Table 2: Identification of different sero- and subtypes of IBDV strains. Determination of the presence of VP5 proteins.

IBDV-serotype	IBDV-subtype	IBDV-strain	R1928	RαVP5	DIE7
I	Classical	D78	+	+	+
I	Classical	228TC	+	+	+
I	Classical	PBG98	+	+	+
I	Classical	Ram0404	+	+	+
I	Classical	IBDV/EK	+	+	+
I	Classical	IBDV/VP5 <sup>-</sup>	+	-	-
I	GLS	GLS	+	+	+
I	Variant-E	8903	+	+	+
II	TY89	TY89	+	+	+

10 From these data it can be concluded that the different strains of IBDV belonging to different sero- and subtypes do express the VP5-gene. Furthermore, the recombinant VP5<sup>-</sup> IBDV vaccine strain can be differentiated from field and vaccine viruses, thereby enabling the recombinant VP5<sup>-</sup> virus to be used as a marker vaccine.

### Example 3

15 In vivo testing of the recombinant VP5<sup>+</sup> and VP5<sup>-</sup> IBDV vaccines in comparison with a commercial available live IBDV vaccine.

20 **Preparation of IBDV vaccine.** Primary chicken embryo fibroblast (CEF) cells were prepared at a final concentration of  $2 \times 10^6$ /ml. The cells were cultured in Eagles minimum essential medium containing 5% fetal calf serum. To 25 ml of this cell suspension 0.1 ml

IBDV/EK or IBDV/VP5<sup>-</sup> virus (having an infectious titre of about 3.0 log<sub>10</sub> TCID<sub>50</sub>/ml) was added. After incubation for 5 days in a high-humidity incubator at 37°C, the total suspension was used in the animal experiment without further purification. The infectious titre of the supernatant was 10<sup>7.1</sup> TCID<sub>50</sub>/ml.

- 5 **Animal experiment.** In this study the potency of different vaccines (VP5 positive strain IBDV/EK and a VP5 negative strain IBDV/VP5<sup>-</sup>, and the commercial available IBDV vaccine Nobilis strain D78, Intervet International B.V., NL) was investigated. SPF chicks of 3 weeks old were treated as indicated in the treatment schedule.

Treatment Schedule:

Days after vaccination	Groups			
	1	2	3	4
00	IBDV/EK	IBDV/VP5 <sup>-</sup>	D78	-
03	x	x1	x	x
07	x,bl	x1,bl	x,b	x,bl
14	x,bl	x,bl	x,bl	x,bl
20	x,bl	x,bl	x,bl	x,bl
21	ch	ch	ch	ch
24	x	x	x	x
31	+	+	+	+

VP5<sup>+</sup> Bursal disease vaccination with VP5 positive vaccine clone, eye-drop route, dose 10<sup>4.6</sup> TCID<sub>50</sub>/animal, 0.1 ml/animal.

VP5<sup>-</sup> Bursal disease vaccination with VP5 negative vaccine clone, eye-drop route, dose 10<sup>5.9</sup> TCID<sub>50</sub>/animal, 0.1 ml/animal.

D78 Bursal disease vaccination with IBDV VACCINE NOBILIS STRAIN D78, eye-drop route, one field dose.

ch Challenge with Bursal disease virus, Farragher strain F52/70, eye-drop route, dose 10<sup>2.0</sup> CID<sub>50</sub>/animal, 0.1 ml/animal.

20 bl Serological examination; VN-test and/or Western blotting.

x Histological examination (H.E. staining) and MCA-8 ELISA on bursae.

- x1 Histological examination (H.E. staining) and MCA-8 ELISA on bursae and reisolation of virus from bursa of Fabricius.
- + Clinical examination and after 10 days histological examination of the bursa.

## 5 **Detection of virus in the bursa of Fabricius.**

Three, 7, 14 and 20 days after eye-drop vaccination, animals were sacrificed and blood and bursae obtained. The presence of virus in the bursa was determined with an enzyme-linked immunosorbent assay (ELISA) making use of the monoclonal antibody 8 (MAB-8). MAB-8 is directed specifically against IBDV. Data are depicted in Table 3.

- 10 Furthermore, 3 and 7 days after vaccination, bursae from animals of group 2 were investigated for the presence of the recombinant VP5<sup>-</sup> virus. For that purpose bursae were homogenised and cultured on chicken embryo fibroblasts. The presence of the VP5<sup>-</sup> virus was determined by IFT using polyclonal rabbit sera against IBDV or VP5 or monoclonal antibodies against VP5. From 13 out of 15 bursae (87%) investigated, VP5<sup>-</sup> virus could be reisolated and
- 15 identified (positive for R1928 and negative for R $\alpha$ VP5 and DIE7). This indicates that the virus upon animal passage is still VP5<sup>-</sup>, indicating that the virus is stable and does not revert to VP5<sup>+</sup>. Furthermore, by using the different poly- and monoclonal antibodies VP5<sup>-</sup> vaccine virus can be discriminated from all other vaccine and/or field IBDV viruses. Therefore, the VP5<sup>-</sup> vaccine may be used as a marker vaccine.

- 20 Three days after challenge no virus could be detected in groups 1, 2 and 3 with the MCA-8 ELISA. In contrast, all animals of group 4 (non-vaccinated control group) contained challenge virus in the bursa of Fabricius, 3 days after challenge. The results show that animals vaccinated with recombinant VP5<sup>+</sup> (group 1), recombinant VP5<sup>-</sup> (group 2) and IBDV vaccine Nobilis D78 (group 3) were protected against severe challenge.

**Table 3:** Individual data for detection of virus in the bursa of Fabricius with the MCA-8 ELISA at different days after vaccination or challenge.

	Days after vaccination→				Days after challenge	
	3	7	14	20	3	
Group↓	Virus detection by ELISA↓					Protection↓
1 VP5 <sup>+</sup>	2/8*	1/7	0/2	0/3	0/5	100%
2 VP5 <sup>-</sup>	0/8	0/7	0/2	0/3	0/5	100%
3 D78	1/8	6/7	0/2	0/3	0/5	100%
4 -	0/8	0/7	0/2	0/3	5/5	0%

5 \*Number of positive bursae per total number tested.

#### Detection of lesions in the bursa of Fabricius.

10 The microscopic average lesion score induced by the different IBDV (recombinant) vaccines or the challenge virus are depicted in Table 4.

Before challenge, animals vaccinated with the recombinant VP5<sup>+</sup> IBDV vaccine (group 1) or vaccinated with IBDV vaccine Nobilis D78 (group 3) showed mild to moderate lesions in the bursa. Three days after challenge only chronic lesions were observed in the bursa of Fabricius, indicating that the animals of groups 1 and 3 were protected against challenge. Furthermore, 10 days after challenge only very mild lesions (0-20% lymphocytic depletion) were observed in the bursa of the animals vaccinated with VP5<sup>+</sup> recombinant IBDV vaccine or with Nobilis vaccine D78. In contrast animals not vaccinated and challenged showed severe lesions 10 days after challenge. In other words all animals (100%) of groups 1 and 3, vaccinated with the VP5<sup>+</sup> recombinant IBDV vaccine or with Nobilis vaccine D78 were protected against severe challenge.

20 Three, 7, 14 and 20 days after vaccination and 3 and 10 days after challenge with the recombinant VP5<sup>-</sup> IBDV vaccine, animals of group 2 showed no to hardly any lesions (0-20% lymphocytic depletion) in the bursa. All animals of group 2, vaccinated with the VP5<sup>-</sup> recombinant IBDV vaccine, were protected against severe challenge. When animals vaccinated with the recombinant VP5<sup>-</sup> IBDV vaccine are compared to animals of groups 1 or 3 (vaccinated



with a recombinant VP5<sup>+</sup> or commercial available vaccine) the recombinant VP5<sup>-</sup> vaccine induces less lesions and therefore, is safer, milder than the vaccines tested in this experiment.

Three days post-challenge, all non-vaccinated animals of group 4 showed severe acute lesions in the bursa (total lymphocyte depletion, score 5.0). Ten days after challenge, all animals ( 17 out of 17 animals) showed total lymphocytic depletion, indicating that these animals were not protected against severe challenge. Animals that died after challenge, all showed severe lesions in the bursa of Fabricius. It was concluded that control group 4 was not protected against severe challenge indicating that the test conditions were optimal.

10 **Table 4:** Average bursal lesion score at different days after vaccination or challenge. The average lesion score is calculated as follows: all lesion scores from the animals per group on a certain day are added. This number is then divided by the total number of animals investigated in that group on that day. Individual scores range from 1 to 5. Score 0 = no lymphocytic depletion, score 1 = 0 - 20%; score 2 = 20 - 40%; score 3 = 40 - 60%; score 4 = 60 - 80% and score 5 = 80 - 100 % lymphocytic depletion (total lymphocytic depletion).

	Days after vaccination→				Days after challenge→		
	3	7	14	20	3	10	
Group↓	Bursal lesions score↓						Protection↓
1 VP5 <sup>+</sup>	0.8	2.9	1.0	1.0	1.0 <sup>c</sup>	0.6	100%
2 VP5 <sup>-</sup>	0.0	0.0	0.5	0.0	0.0 <sup>c</sup>	0.1	100%
3 D78	0.1	2.4	3.5	2.0	2.8 <sup>c</sup>	1.1	100%
4 -	0.0	0.0	0.0	0.0	5.0 <sup>a</sup>	5.0	0%

<sup>a</sup> Acute lesions <sup>c</sup> Chronic lesions

## 20 Serological response.

The serological response of the animals was determined by measuring the ability of blood serum to neutralise a classical infectious bursal disease virus strain in a virus neutralising (VN) test. Serum was investigated 3, 7, 14 and 20 days after vaccination. The average neutralising titres are shown in Table 5.

The results show that recombinant IBDV vaccine VP5<sup>+</sup> applied to chickens of group 1 induced a good and high serological response 20 days after vaccination which is comparable to the serological response of the chickens vaccinated with the commercial IBDV vaccine Nobilis strain D78 (group 3). The recombinant IBDV vaccine VP5<sup>-</sup> applied to chickens of group 2 induced also a good serological response. A titre of 9.4 log<sub>2</sub> was observed 20 days after vaccination. The serological response induced by the recombinant VP5<sup>-</sup> IBDV vaccine was delayed when compared to the serological response induced by the recombinant IBDV VP5<sup>+</sup> vaccine or the commercial IBDV vaccine Nobilis strain D78.

The non-vaccinated group 4 showed no serological response to IBDV.

**Table 5:** Average IBDV-VN-titres for groups 1 to 4 at different days after vaccination, expressed as log<sub>2</sub> of the dilution.

Group	Days after vaccination			
	3	7	14	20
1 VP5 <sup>+</sup>	$\leq 1.0 \pm 0.0$	$7.1 \pm 1.7$	$10.2 \pm 1.4$	$11.9 \pm 1.8$
2 VP5 <sup>-</sup>	$\leq 1.0 \pm 0.0$	$2.1 \pm 1.7$	$6.3 \pm 2.9$	$9.4 \pm 1.4$
3 D78	$\leq 1.0 \pm 0.0$	$5.2 \pm 2.8$	$10.3 \pm 1.3$	$11.6 \pm 1.5$
4 -	$\leq 1.0 \pm 0.0$	$\leq 1.0 \pm 0.0$	$\leq 1.0 \pm 0.0$	$\leq 1.0 \pm 0.0$

#### **Serological differentiation between antisera.**

The serological response against VP5 was investigated by making use of western blot analysis. For this purpose the VP5 protein was expressed in the E. coli or baculo expression system. The expressed proteins were separated by SDS PAGE. Next the proteins were electroblotted onto a nitro-cellulose membrane. Thereafter, the membrane was cut into lanes and the lanes were incubated with rabbit anti-VP5 serum, chicken serum directed against VP5<sup>+</sup> recombinant vaccine, chicken serum directed against VP5<sup>-</sup> recombinant vaccine or negative serum from SPF chickens. Data are summarised in Table 6. As can be seen from Table 6, the VP5<sup>-</sup> serum does not induce a serological response against VP5. In contrast the rabbit anti-VP5 serum and chicken serum directed against VP5<sup>+</sup> recombinant vaccine do recognise the VP5-

protein and thus induces a serological response against VP5. This indicates that chicken serum may be used to investigate if animals are exposed to a virus that expresses the VP5 protein (e.g. field virus) or to the VP5<sup>-</sup> recombinant vaccine.

- 5      Table 6: Western blot analysis. Serum from animals vaccinated with VP5<sup>+</sup> or VP5-recombinant vaccine as well as SPF chicken serum and anti VP5-rabbit serum were investigated for their reaction with the VP5-protein.

Identification of serum sample	Immuno-blot
VP5 <sup>+</sup> vaccinated animal, serum sample 20d after vaccination	positive
VP5 <sup>-</sup> vaccinated animal, serum sample 20d after vaccination	negative
Non-vaccinated control, serum sample at 20d	negative
Rabbit anti VP5 serum	positive

10

#### **Mortality and clinical signs.**

None of the animals vaccinated with VP5<sup>+</sup> IBDV vaccine (group 1), vaccinated with recombinant VP5<sup>-</sup> IBDV vaccine (group 2) or vaccinated with the commercial IBDV vaccine

15      Nobilis strain D78 (group 3), died or showed clinical signs of infectious bursal disease after challenge, indicating that the animals were protected against severe challenge. All animals in the non-vaccinated control group were not protected against severe challenge.

20

### LEGENDS TO THE FIGURES

#### Figure 1

Genomic organization of segment A and segment B of IBDV. The numbers indicate the nucleotide positions of the start, end and coding region on the segments.

#### Figure 2

Construction of genomic cDNA clones for the preparation of IBDV/VP5<sup>-</sup>. Plasmid pAD78/EK contains the complete D78 segment A cDNA encoding the polyprotein (VP2-VP4-VP3) and VP5. Plasmid pBP2 contains the complete strain P2 segment B encoding VP1. Mutations were introduced in plasmid pAD78/VP5<sup>-</sup> altering the methionine start codon for VP5 into arginine and creating an artificial Afl II cleavage site. Recombinant plasmids were linearized with the underlined restriction enzymes, followed by T7 polymerase transcription.

#### Figure 3

Construction of genomic cDNA clones for the preparation of IBDV/VP5<sup>-</sup>-2. Plasmid pAD78/EK contains the complete D78 segment A cDNA encoding the polyprotein (VP2-VP4-VP3) and VP5. Plasmid pBD78 contains the complete strain D78 segment B encoding VP1. Mutations were introduced in plasmid pAD78/VP5<sup>-</sup> altering the methionine start codon for VP5 into glutamic acid and creating an artificial BstBI cleavage site. Further mutations were introduced in the arginine and glutamine codon. Recombinant plasmids were linearized with the underlined restriction enzymes, followed by T7 polymerase transcription.

#### Figure 4

Radioimmunoprecipitation of proteins from CEC infected cells with recombinant IBDV. CEC infected cells with IBDV/VP5<sup>-</sup> (lanes 1-3), IBDV/EK (lanes 4-6) and uninfected controls were immunoprecipitated with rabbit anti-IBDV serum (lanes 1, 4, 7), rabbit anti-VP5 serum (lanes 2, 5, 8) and mAb DIE 7 (lanes 3, 6, 9). Position of molecular mass markers (M) is indicated. Location of the viral proteins VP2, VP3, VP4 and VP5 are marked.

#### Figure 5

Replication kinetics of IBDV/EK and IBDV/VP5<sup>-</sup>. Infectious titers of supernatants (vertical axis) are determined at the times indicated.

## SEQUENCE LISTING

## 5 (1) GENERAL INFORMATION:

## (i) APPLICANT:

- (A) NAME: Azko Nobel N.V.  
(B) STREET: Velperweg 76  
10 (C) CITY: Arnhem  
(E) COUNTRY: The Netherlands  
(F) POSTAL CODE (ZIP): 6824 BM  
(G) TELEPHONE: 0412 666379  
15 (H) TELEFAX: 0412 650592

(ii) TITLE OF INVENTION: Recombinant birnavirus vaccine

(iii) NUMBER OF SEQUENCES: 8

## 20 (iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk  
(B) COMPUTER: IBM PC compatible  
(C) OPERATING SYSTEM: PC-DOS/MS-DOS  
25 (D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

## (2) INFORMATION FOR SEQ ID NO: 1:

## 30 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2827 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

35 (ii) MOLECULE TYPE: cDNA

## (ix) FEATURE:

- (A) NAME/KEY: CDS  
40 (B) LOCATION: 112..2745

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

45 GGATACGATG GGTCTGACCC TCTGGGAGTC ACGAATTAAC GTGGCTACTA GGGGCGATAC

60

	CCGCCGCTGG CCGCCACGTT AGTGGCTCCT CTTCTTGATG ATTCTGCCAC C ATG AGT	117
	Met Ser	
	1	
5	GAC ATT TTC AAC AGT CCA CAG GCG CGA AGC ACG ATC TCA GCA GCG TTC	165
	Asp Ile Phe Asn Ser Pro Gln Ala Arg Ser Thr Ile Ser Ala Ala Phe	
	5 10 15	
10	GGC ATA AAG CCT ACT GCT GGA CAA GAC GTG GAA GAA CTC TTG ATC CCT	213
	Gly Ile Lys Pro Thr Ala Gly Gln Asp Val Glu Glu Leu Leu Ile Pro	
	20 25 30	
15	AAA GTT TGG GTG CCA CCT GAG GAT CCG CTT GCC AGC CCT AGT CGA CTG	261
	Lys Val Trp Val Pro Pro Glu Asp Pro Leu Ala Ser Pro Ser Arg Leu	
	35 40 45 50	
20	GCA AAG TTC CTC AGA GAG AAC GGC TAC AAA GTT TTG CAG CCA CGG TCT	309
	Ala Lys Phe Leu Arg Glu Asn Gly Tyr Lys Val Leu Gln Pro Arg Ser	
	55 60 65	
25	CTG CCC GAG AAT GAG GAG TAT GAG ACC GAC CAA ATA CTC CCA GAC TTA	357
	Leu Pro Glu Asn Glu Glu Tyr Glu Thr Asp Gln Ile Leu Pro Asp Leu	
	70 75 80	
30	GCA TGG ATG CGA CAG ATA GAA GGG GCT GTT TTA AAA CCC ACT CTA TCT	405
	Ala Trp Met Arg Gln Ile Glu Gly Ala Val Leu Lys Pro Thr Leu Ser	
	85 90 95	
35	CTC CCT ATT GGA GAT CAG GAG TAC TTC CCA AAG TAC TAC CCA ACA CAT	453
	Leu Pro Ile Gly Asp Gln Glu Tyr Phe Pro Lys Tyr Tyr Pro Thr His	
	100 105 110	
40	CGC CCT AGC AAG GAG AAG CCC AAT GCG TAC CCG CCA GAC ATC GCA CTA	501
	Arg Pro Ser Lys Glu Lys Pro Asn Ala Tyr Pro Pro Asp Ile Ala Leu	
	115 120 125 130	
45	CTC AAG CAG ATG ATT TAC CTG TTT CTC CAG GTT CCA GAG GCC AAC GAG	549
	Leu Lys Gln Met Ile Tyr Leu Phe Leu Gln Val Pro Glu Ala Asn Glu	
	135 140 145	
50	GGC CTA AAG GAT GAA GTA ACC CTC TTG ACC CAA AAC ATA AGG GAC AAG	597
	Gly Leu Lys Asp Glu Val Thr Leu Leu Thr Gln Asn Ile Arg Asp Lys	
	150 155 160	
55	GCC TAT GGA AGT GGG ACC TAC ATG GGA CAA GCA AAT CGA CTT GTG GCC	645
	Ala Tyr Gly Ser Gly Thr Tyr Met Gly Gln Ala Asn Arg Leu Val Ala	
	165 170 175	

	ATG AAG GAG GTC GCC ACT GGA AGA AAC CCA AAC AAG GAT CCT CTA AAG	693
	Met Lys Glu Val Ala Thr Gly Arg Asn Pro Asn Lys Asp Pro Leu Lys	
	180 185 190	
5	CTT GGG TAC ACT TTT GAG AGC ATC GCG CAG CTA CTT GAC ATC ACA CTA	741
	Leu Gly Tyr Thr Phe Glu Ser Ile Ala Gln Leu Leu Asp Ile Thr Leu	
	195 200 205 210	
10	CCG GTA GGC CCA CCC GGT GAG GAT GAC AAG CCC TGG GTG CCA CTC ACA	789
	Pro Val Gly Pro Pro Gly Glu Asp Asp Lys Pro Trp Val Pro Leu Thr	
	215 220 225	
15	AGA GTG CCG TCA CGG ATG TTG GTG CTG ACG GGA GAC GTA GAT GGC GAC	837
	Arg Val Pro Ser Arg Met Leu Val Leu Thr Gly Asp Val Asp Gly Asp	
	230 235 240	
20	TTT GAG GTT GAA GAT TAC CTT CCC AAA ATC AAC CTC AAG TCA TCA AGT	885
	Phe Glu Val Glu Asp Tyr Leu Pro Lys Ile Asn Leu Lys Ser Ser Ser	
	245 250 255	
	GGA CTA CCA TAT GTA GGT CGC ACC AAA GGA GAG ACA ATT GGC GAG ATG	933
	Gly Leu Pro Tyr Val Gly Arg Thr Lys Gly Glu Thr Ile Gly Glu Met	
	260 265 270	
25	ATA GCT ATC TCA AAC CAG TTT CTC AGA GAG CTA TCA ACA CTG TTG AAG	981
	Ile Ala Ile Ser Asn Gln Phe Leu Arg Glu Leu Ser Thr Leu Leu Lys	
	275 280 285 290	
30	CAA GGT GCA GGG ACA AAG GGG TCA AAC AAG AAG AAG CTA CTC AGC ATG	1029
	Gln Gly Ala Gly Thr Lys Gly Ser Asn Lys Lys Lys Leu Leu Ser Met	
	295 300 305	
35	TTA AGT GAC TAT TGG TAC TTA TCA TGC GGG CTT TTG TTT CCA AAG GCT	1077
	Leu Ser Asp Tyr Trp Tyr Leu Ser Cys Gly Leu Leu Phe Pro Lys Ala	
	310 315 320	
40	GAA AGG TAC GAC AAA AGT ACA TGG CTC ACC AAG ACC CGG AAC ATA TGG	1125
	Glu Arg Tyr Asp Lys Ser Thr Trp Leu Thr Lys Thr Arg Asn Ile Trp	
	325 330 335	
	TCA GCT CCA TCC CCA ACA CAC CTC ATG ATC TCT ATG ATC ACC TGG CCC	1173
	Ser Ala Pro Ser Pro Thr His Leu Met Ile Ser Met Ile Thr Trp Pro	
	340 345 350	
45	GTG ATG TCC AAC AGC CCA AAT AAC GTG TTG AAC ATT GAA GGG TGT CCA	1221
	Val Met Ser Asn Ser Pro Asn Asn Val Leu Asn Ile Glu Gly Cys Pro	
	355 360 365 370	

	TCA CTC TAC AAA TTC AAC CCG TTC AGA GGA GGG TTG AAC AGG ATC GTC	1269
	Ser Leu Tyr Lys Phe Asn Pro Phe Arg Gly Gly Leu Asn Arg Ile Val	
	375 380 385	
5	GAG TGG ATA TTG GCC CCG GAA GAA CCC AAG GCT CTT GTA TAT GCG GAC	1317
	Glu Trp Ile Leu Ala Pro Glu Glu Pro Lys Ala Leu Val Tyr Ala Asp	
	390 395 400	
10	AAC ATA TAC ATT GTC CAC TCA AAC ACG TGG TAC TCA ATT GAC CTA GAG	1365
	Asn Ile Tyr Ile Val His Ser Asn Thr Trp Tyr Ser Ile Asp Leu Glu	
	405 410 415	
15	AAG GGT GAG GCA AAC TGC ACT CGC CAA CAC ATG CAA GCC GCA ATG TAC	1413
	Lys Gly Glu Ala Asn Cys Thr Arg Gln His Met Gln Ala Ala Met Tyr	
	420 425 430	
20	TAC ATA CTC ACC AGA GGG TGG TCA GAC AAC GGC GAC CCA ATG TTC AAT	1461
	Tyr Ile Leu Thr Arg Gly Trp Ser Asp Asn Gly Asp Pro Met Phe Asn	
	435 440 445 450	
	CAA ACA TGG GCC ACC TTT GCC ATG AAC ATT GCC CCT GCT CTA GTG GTG	1509
	Gln Thr Trp Ala Thr Phe Ala Met Asn Ile Ala Pro Ala Leu Val Val	
	455 460 465	
25	GAC TCA TCG TGC CTG ATA ATG AAC CTG CAA ATT AAG ACC TAT GGT CAA	1557
	Asp Ser Ser Cys Leu Ile Met Asn Leu Gln Ile Lys Thr Tyr Gly Gln	
	470 475 480	
30	GGC AGC GGG AAT GCA GCC ACG TTC ATC AAC AAC CAC CTC TTG AGC ACA	1605
	Gly Ser Gly Asn Ala Ala Thr Phe Ile Asn Asn His Leu Leu Ser Thr	
	485 490 495	
35	CTA GTG CTT GAC CAG TGG AAC CTG ATG AGA CAG CCC AGA CCA GAC AGC	1653
	Leu Val Leu Asp Gln Trp Asn Leu Met Arg Gln Pro Arg Pro Asp Ser	
	500 505 510	
40	GAG GAG TTC AAA TCA ATT GAG GAC AAG CTA GGT ATC AAC TTT AAG ATT	1701
	Glu Glu Phe Lys Ser Ile Glu Asp Lys Leu Gly Ile Asn Phe Lys Ile	
	515 520 525 530	
	GAG AGG TCC ATT GAT GAT ATC AGG GGC AAG CTG AGA CAG CTT GTC CTC	1749
	Glu Arg Ser Ile Asp Asp Ile Arg Gly Lys Leu Arg Gln Leu Val Leu	
	535 540 545	
45	CTT GCA CAA CCA GGG TAC CTG AGT GGG GGG GTT GAA CCA GAA CAA TCC	1797
	Leu Ala Gln Pro Gly Tyr Leu Ser Gly Gly Val Glu Pro Glu Gln Ser	
	550 555 560	



	AGC CCA ACT GTT GAG CTT GAC CTA CTA GGG TGG TCA GCT ACA TAC AGC Ser Pro Thr Val Glu Leu Asp Leu Leu Gly Trp Ser Ala Thr Tyr Ser	1845
	565 570 575	
5	AAA GAT CTC GGG ATC TAT GTG CCG GTG CTT GAC AAG GAA CGC CTA TTT Lys Asp Leu Gly Ile Tyr Val Pro Val Leu Asp Lys Glu Arg Leu Phe	1893
	580 585 590	
10	TGT TCT GCT GCG TAT CCC AAG GGA GTA GAG AAC AAG AGT CTC AAG TCC Cys Ser Ala Ala Tyr Pro Lys Gly Val Glu Asn Lys Ser Leu Lys Ser	1941
	595 600 605 610	
15	AAA GTC GGG ATC GAG CAG GCA TAC AAG GTA GTC AGG TAT GAG GCG TTG Lys Val Gly Ile Glu Gln Ala Tyr Lys Val Val Arg Tyr Glu Ala Leu	1989
	615 620 625	
20	AGG TTG GTA GGT GGT TGG AAC TAC CCA CTC CTG AAC AAA GCC TGC AAG Arg Leu Val Gly Gly Trp Asn Tyr Pro Leu Leu Asn Lys Ala Cys Lys	2037
	630 635 640	
	AAT AAC GCA GGC GCC GCT CGG CGG CAT CTG GAG GCC AAG GGG TTC CCA Asn Asn Ala Gly Ala Ala Arg Arg His Leu Glu Ala Lys Gly Phe Pro	2085
	645 650 655	
25	CTC GAC GAG TTC CTA GCC GAG TGG TCT GAG CTG TCA GAG TTC GGT GAG Leu Asp Glu Phe Leu Ala Glu Trp Ser Glu Leu Ser Glu Phe Gly Glu	2133
	660 665 670	
30	GCC TTC GAA GGC TTC AAT ATC AAG CTG ACC GTA ACA TCT GAG AGC CTA Ala Phe Glu Gly Phe Asn Ile Lys Leu Thr Val Thr Ser Glu Ser Leu	2181
	675 680 685 690	
35	GCC GAA CTG AAC AAG CCA GTA CCC CCC AAG CCC CCA AAT GTC AAC AGA Ala Glu Leu Asn Lys Pro Val Pro Pro Lys Pro Pro Asn Val Asn Arg	2229
	695 700 705	
40	CCA GTC AAC ACT GGG GGA CTC AAG GCA GTC AGC AAC GCC CTC AAG ACC Pro Val Asn Thr Gly Gly Leu Lys Ala Val Ser Asn Ala Leu Lys Thr	2277
	710 715 720	
	GGT CGG TAC AGG AAC GAA GCC GGA CTG AGT GGT CTC GTC CTT CTA GCC Gly Arg Tyr Arg Asn Glu Ala Gly Leu Ser Gly Leu Val Leu Leu Ala	2325
	725 730 735	
45	ACA GCA AGA AGC CGT CTG CAA GAT GCA GTT AAG GCC AAG GCA GAA GCC Thr Ala Arg Ser Arg Leu Gln Asp Ala Val Lys Ala Lys Ala Glu Ala	2373
	740 745 750	



45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

	Met	Ser	Asp	Ile	Phe	Asn	Ser	Pro	Gln	Ala	Arg	Ser	Thr	Ile	Ser	Ala	
	1				5					10					15		
5	Ala	Phe	Gly	Ile	Lys	Pro	Thr	Ala	Gly	Gln	Asp	Val	Glu	Glu	Leu	Leu	
				20					25					30			
	Ile	Pro	Lys	Val	Trp	Val	Pro	Pro	Glu	Asp	Pro	Leu	Ala	Ser	Pro	Ser	
			35					40					45				
10	Arg	Leu	Ala	Lys	Phe	Leu	Arg	Glu	Asn	Gly	Tyr	Lys	Val	Leu	Gln	Pro	
		50					55					60					
	Arg	Ser	Leu	Pro	Glu	Asn	Glu	Glu	Tyr	Glu	Thr	Asp	Gln	Ile	Leu	Pro	
15		65				70					75				80		
	Asp	Leu	Ala	Trp	Met	Arg	Gln	Ile	Glu	Gly	Ala	Val	Leu	Lys	Pro	Thr	
				85					90					95			
20	Leu	Ser	Leu	Pro	Ile	Gly	Asp	Gln	Glu	Tyr	Phe	Pro	Lys	Tyr	Tyr	Pro	
			100						105					110			
	Thr	His	Arg	Pro	Ser	Lys	Glu	Lys	Pro	Asn	Ala	Tyr	Pro	Pro	Asp	Ile	
			115					120					125				
25	Ala	Leu	Leu	Lys	Gln	Met	Ile	Tyr	Leu	Phe	Leu	Gln	Val	Pro	Glu	Ala	
		130					135					140					
	Asn	Glu	Gly	Leu	Lys	Asp	Glu	Val	Thr	Leu	Leu	Thr	Gln	Asn	Ile	Arg	
30		145				150					155				160		
	Asp	Lys	Ala	Tyr	Gly	Ser	Gly	Thr	Tyr	Met	Gly	Gln	Ala	Asn	Arg	Leu	
				165					170					175			
35	Val	Ala	Met	Lys	Glu	Val	Ala	Thr	Gly	Arg	Asn	Pro	Asn	Lys	Asp	Pro	
			180						185					190			
	Leu	Lys	Leu	Gly	Tyr	Thr	Phe	Glu	Ser	Ile	Ala	Gln	Leu	Leu	Asp	Ile	
			195					200					205				
40	Thr	Leu	Pro	Val	Gly	Pro	Pro	Gly	Glu	Asp	Asp	Lys	Pro	Trp	Val	Pro	
		210					215					220					
	Leu	Thr	Arg	Val	Pro	Ser	Arg	Met	Leu	Val	Leu	Thr	Gly	Asp	Val	Asp	
45		225				230				235				240			
	Gly	Asp	Phe	Glu	Val	Glu	Asp	Tyr	Leu	Pro	Lys	Ile	Asn	Leu	Lys	Ser	
				245					250					255			

Ser Ser Gly Leu Pro Tyr Val Gly Arg Thr Lys Gly Glu Thr Ile Gly  
 260 265 270

5 Glu Met Ile Ala Ile Ser Asn Gln Phe Leu Arg Glu Leu Ser Thr Leu  
 275 280 285

Leu Lys Gln Gly Ala Gly Thr Lys Gly Ser Asn Lys Lys Lys Leu Leu  
 290 295 300

10 Ser Met Leu Ser Asp Tyr Trp Tyr Leu Ser Cys Gly Leu Leu Phe Pro  
 305 310 315 320

Lys Ala Glu Arg Tyr Asp Lys Ser Thr Trp Leu Thr Lys Thr Arg Asn  
 325 330 335

15 Ile Trp Ser Ala Pro Ser Pro Thr His Leu Met Ile Ser Met Ile Thr  
 340 345 350

Trp Pro Val Met Ser Asn Ser Pro Asn Asn Val Leu Asn Ile Glu Gly  
 20 355 360 365

Cys Pro Ser Leu Tyr Lys Phe Asn Pro Phe Arg Gly Gly Leu Asn Arg  
 370 375 380

25 Ile Val Glu Trp Ile Leu Ala Pro Glu Glu Pro Lys Ala Leu Val Tyr  
 385 390 395 400

Ala Asp Asn Ile Tyr Ile Val His Ser Asn Thr Trp Tyr Ser Ile Asp  
 405 410 415

30 Leu Glu Lys Gly Glu Ala Asn Cys Thr Arg Gln His Met Gln Ala Ala  
 420 425 430

Met Tyr Tyr Ile Leu Thr Arg Gly Trp Ser Asp Asn Gly Asp Pro Met  
 35 435 440 445

Phe Asn Gln Thr Trp Ala Thr Phe Ala Met Asn Ile Ala Pro Ala Leu  
 450 455 460

40 Val Val Asp Ser Ser Cys Leu Ile Met Asn Leu Gln Ile Lys Thr Tyr  
 465 470 475 480

Gly Gln Gly Ser Gly Asn Ala Ala Thr Phe Ile Asn Asn His Leu Leu  
 485 490 495

45 Ser Thr Leu Val Leu Asp Gln Trp Asn Leu Met Arg Gln Pro Arg Pro  
 500 505 510

	Asp Ser Glu Glu Phe Lys Ser Ile Glu Asp Lys Leu Gly Ile Asn Phe	
	515	520 525
5	Lys Ile Glu Arg Ser Ile Asp Asp Ile Arg Gly Lys Leu Arg Gln Leu	
	530	535 540
	Val Leu Leu Ala Gln Pro Gly Tyr Leu Ser Gly Gly Val Glu Pro Glu	
	545	550 555 560
10	Gln Ser Ser Pro Thr Val Glu Leu Asp Leu Leu Gly Trp Ser Ala Thr	
	565	570 575
	Tyr Ser Lys Asp Leu Gly Ile Tyr Val Pro Val Leu Asp Lys Glu Arg	
	580	585 590
15	Leu Phe Cys Ser Ala Ala Tyr Pro Lys Gly Val Glu Asn Lys Ser Leu	
	595	600 605
	Lys Ser Lys Val Gly Ile Glu Gln Ala Tyr Lys Val Val Arg Tyr Glu	
20	610	615 620
	Ala Leu Arg Leu Val Gly Gly Trp Asn Tyr Pro Leu Leu Asn Lys Ala	
	625	630 635 640
25	Cys Lys Asn Asn Ala Gly Ala Ala Arg Arg His Leu Glu Ala Lys Gly	
	645	650 655
	Phe Pro Leu Asp Glu Phe Leu Ala Glu Trp Ser Glu Leu Ser Glu Phe	
	660	665 670
30	Gly Glu Ala Phe Glu Gly Phe Asn Ile Lys Leu Thr Val Thr Ser Glu	
	675	680 685
	Ser Leu Ala Glu Leu Asn Lys Pro Val Pro Pro Lys Pro Pro Asn Val	
35	690	695 700
	Asn Arg Pro Val Asn Thr Gly Gly Leu Lys Ala Val Ser Asn Ala Leu	
	705	710 715 720
40	Lys Thr Gly Arg Tyr Arg Asn Glu Ala Gly Leu Ser Gly Leu Val Leu	
	725	730 735
	Leu Ala Thr Ala Arg Ser Arg Leu Gln Asp Ala Val Lys Ala Lys Ala	
	740	745 750
45	Glu Ala Glu Lys Leu His Lys Ser Lys Pro Asp Asp Pro Asp Ala Asp	
	755	760 765

Trp Phe Glu Arg Ser Glu Thr Leu Ser Asp Leu Leu Glu Lys Ala Asp  
 770 775 780

5 Ile Ala Ser Lys Val Ala His Ser Ala Leu Val Glu Thr Ser Asp Ala  
 785 790 795 800

Leu Glu Ala Val Gln Ser Thr Ser Val Tyr Thr Pro Lys Tyr Pro Glu  
 805 810 815

10 Val Lys Asn Pro Gln Thr Ala Ser Asn Pro Val Val Gly Leu His Leu  
 820 825 830

Pro Ala Lys Arg Ala Thr Gly Val Gln Ala Ala Leu Leu Gly Ala Gly  
 835 840 845

15 Thr Ser Arg Pro Met Gly Met Glu Ala Pro Thr Arg Ser Lys Asn Ala  
 850 855 860

20 Val Lys Met Ala Lys Arg Arg Gln Arg Gln Lys Glu Ser Arg  
 865 870 875

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 3261 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 97..531

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

GGATACGATC GGTCTGACCC CGGGGGAGTC ACCCGGGGAC AGGCCGTCAA GGCCTTGTTT 60

45 CAGGATGGGA CTCCTCCTTC TACAACGCTA TCATTG ATG GTT AGT AGA GAT CAG 114  
 Met Val Ser Arg Asp Gln

	ACA AAC GAT CGC AGC GAT GAC AAA CCT GCA AGA TCA AAC CCA ACA GAT	162
	Thr Asn Asp Arg Ser Asp Asp Lys Pro Ala Arg Ser Asn Pro Thr Asp	
	10 15 20	
5	TGT TCC GTT CAT ACG GAG CCT TCT GAT GCC AAC AAC CGG ACC GGC GTC	210
	Cys Ser Val His Thr Glu Pro Ser Asp Ala Asn Asn Arg Thr Gly Val	
	25 30 35	
10	CAT TCC GGA CGA CAC CCT GGA GAA GCA CAC TCT CAG GTC AGA GAC CTC	258
	His Ser Gly Arg His Pro Gly Glu Ala His Ser Gln Val Arg Asp Leu	
	40 45 50	
15	GAC CTA CAA TTT GAC TGT GGG GGA CAC AGG GTC AGG GCT AAT TGT CTT	306
	Asp Leu Gln Phe Asp Cys Gly Gly His Arg Val Arg Ala Asn Cys Leu	
	55 60 65 70	
20	TTT CCC TGG ATT CCC TGG CTC AAT TGT GGG TGC TCA CTA CAC ACT GCA	354
	Phe Pro Trp Ile Pro Trp Leu Asn Cys Gly Cys Ser Leu His Thr Ala	
	75 80 85	
25	GGG CAA TGG GAA CTA CAA GTT CGA TCA GAT GCT CCT GAC TGC CCA GAA	402
	Gly Gln Trp Glu Leu Gln Val Arg Ser Asp Ala Pro Asp Cys Pro Glu	
	90 95 100	
30	CCT ACC GGC CAG TTA CAA CTA CTG CAG GCT AGT GAG TCG GAG TCT CAC	450
	Pro Thr Gly Gln Leu Gln Leu Leu Gln Ala Ser Glu Ser Glu Ser His	
	105 110 115	
35	AGT GAG GTC AAG CAC ACT TCC TGG TGG CGT TTA TGC ACT AAA CGG CAC	498
	Ser Glu Val Lys His Thr Ser Trp Trp Arg Leu Cys Thr Lys Arg His	
	120 125 130	
40	CAT AAA CGC CGT GAC CTT CCA AGG AAG CCT GAG TGA ACT GACA GAT GTT AGCT	551
	His Lys Arg Arg Asp Leu Pro Arg Lys Pro Glu	
	135 140 145	
45	ACAATGGGTT GATGTCTGCA ACAGCCAACA TCAACGACAA AATTGGGAAC GTCCTAGTAG	611
	GGGAAGGGGT CACCGTCCTC AGCTTACCCA CATCATATGA TCTTGGGTAT GTGAGGCTTG	671
	GTGACCCCAT TCCCGCAATA GGGCTTGACC CAAAAATGGT AGCCACATGT GACAGCAGTG	731
	ACAGGCCAG AGTCTACACC ATA ACTGCAG CCGATGATTA CCAATTCTCA TCACAGTACC	791
	AACCAGGTGG GGTAACAATC AACTGTTCT CAGCCAACAT TGATGCCATC ACAAGCCTCA	851
	GCGTTGGGGG AGAGCTCGTG TTTCAAACAA GCGTCCACGG CCTTGTACTG GGCGCCACCA	911

	TCTACCTCAT	AGGCTTTGAT	GGGACAACGG	TAATCACCAG	GGCTGTGGCC	GCAAACAATG	971
	GGCTGACGAC	CGGCACCGAC	AACCTTATGC	CATTCAATCT	TGTGATTCCA	ACAAACGAGA	1031
5	TAACCCAGCC	AATCACATCC	ATCAAAGTGG	AGATAGTGAC	CTCCAAAAGT	GGTGGTCAGG	1091
	CAGGGGATCA	GATGTCATGG	TCGGCAAGAG	GGAGCCTAGC	AGTGACGATC	CATGGTGGCA	1151
	ACTATCCAGG	GGCCCTCCGT	CCCGTCACGC	TAGTGGCCTA	CGAAAGAGTG	GCAACAGGAT	1211
10	CCGTCGTTAC	GGTCGCTGGG	GTGAGCAACT	TCGAGCTGAT	CCCAAATCCT	GAAGTAGCAA	1271
	AGAACCTGGT	TACAGAATAC	GGCCGATTTG	ACCCAGGAGC	CATGAACTAC	ACAAAATTGA	1331
15	TACTGAGTGA	GAGGGACCGT	CTTGGCATCA	AGACCGTCTG	GCCAACAAGG	GAGTACACTG	1391
	ACTTTCGTGA	ATACTTCATG	GAGGTGGCCG	ACCTCAACTC	TCCCCTGAAG	ATTGCAGGAG	1451
	CATTGCGCTT	CAAAGACATA	ATCCGGGCCA	TAAGGAGGAT	AGCTGTGCCG	GTGGTCTCCA	1511
20	CATTGTTCCC	ACCTGCCGCT	CCCCTAGCCC	ATGCAATTGG	GGAAGGTGTA	GACTACCTGC	1571
	TGGGCGATGA	GGCACAGGCT	GCTTCAGGAA	CTGCTCGAGC	CGCGTCAGGA	AAAGCAAGAG	1631
25	CTGCCTCAGG	CCGCATAAGG	CAGCTGACTC	TCGCCGCCGA	CAAGGGGTAC	GAGGTAGTCG	1691
	CGAATCTATT	CCAGGTGCCC	CAGAATCCCG	TAGTCGACGG	GATTCTTGCT	TCACCTGGGG	1751
	TACTCCGCGG	TGCACACAAC	CTCGACTGCG	TGTTAAGAGA	GGGTGCCACG	CTATTCCCTG	1811
30	TGGTTATTAC	GACAGTGGAA	GACGCCATGA	CACCCAAAGC	ATTGAACAGC	AAAATGTTTG	1871
	CTGTCATTGA	AGGCGTGCGA	GAAGACCTCC	AACCTCCATC	TCAAAGAGGA	TCCTTCATAC	1931
35	GAACTCTCTC	TGGACACAGA	GTCTATGGAT	ATGCTCCAGA	TGGGGTACTT	CCACTGGAGA	1991
	CTGGGAGAGA	CTACACCGTT	GTCCCAATAG	ATGATGTCTG	GGACGACAGC	ATTATGCTGT	2051
	CCAAAGATCC	CATACCTCCT	ATTGTGGGAA	ACAGTGGAAG	TCTAGCCATA	GCTTACATGG	2111
40	ATGTGTTTCG	ACCCAAAGTC	CCAATCCATG	TGGCTATGAC	GGGAGCCCTC	AATGCTTGTC	2171
	GCGAGATTGA	GAAAGTAAGC	TTTAGAAGCA	CCAAGCTCGC	CACTGCACAC	CGACTTGGCC	2231
45	TTAGGTTGGC	TGGTCCCGGA	GCATTCGATG	TAAACACCGG	GCCCAACTGG	GCAACGTTCA	2291
	TCAAACGTTT	CCCTCACAAT	CCACGCGACT	GGGACAGGCT	CCCCTACCTC	AACCTACCAT	2351



ACCTTCCACC CAATGCAGGA CGCCAGTACC ACCTTGCCAT GGCTGCATCA GAGTTCAAAG 2411  
 AGACCCCCGA ACTCGAGAGT GCCGTCAGAG CAATGGAAGC AGCAGCCAAC GTGGACCCAC 2471  
 5 TATTCCAATC TGCACTCAGT GTGTTTCATGT GGCTGGAAGA GAATGGGATT GTGACTGACA 2531  
 TGGCCAACTT CGCACTCAGC GACCCGAACG CCCATCGGAT GCGAAATTTT CTTGCAAACG 2591  
 10 CACCACAAGC AGGCAGCAAG TCGCAAAGGG CCAAGTACGG GACAGCAGGC TACGGAGTGG 2651  
 AGGCTCGGGG CCCCACACCA GAGGAAGCAC AGAGGGAAAA AGACACACGG ATCTCAAAGA 2711  
 AGATGGAGAC CATGGGCATC TACTTTGCAA CACCAGAATG GGTAGCACTC AATGGGCACC 2771  
 15 GAGGGCCAAG CCCC GGCCAG CTAAAGTACT GGCAGAACAC ACGAGAAATA CCGGACCCAA 2831  
 ACGAGGACTA TCTAGACTAC GTGCATGCAG AGAAGAGCCG GTTGGCATCA GAAGAACAAA 2891  
 TCCTAAGGGC AGCTACGTCG ATCTACGGGG CTCCAGGACA GGCAGAGCCA CCCCAAGCTT 2951  
 20 TCATAGACGA AGTTGCCAAA GTCTATGAAA TCAACCATGG ACGTGGCCCA AACCAAGAAC 3011  
 AGATGAAAGA TCTGCTCTTG ACTGCGATGG AGATGAAGCA TCGCAATCCC AGGCGGGCTC 3071  
 25 TACCAAAGCC CAAGCCAAAA CCAATGCTC CAACACAGAG ACCCCCTGGT CGGCTGGGCC 3131  
 GCTGGATCAG GACCGTCTCT GATGAGGACC TTGAGTGAGG CTCCTGGGAG TCTCCCGACA 3191  
 CCACCCGCGC AGGTGTGGAC ACCAATTCGG CCTTACAACA TCCCAAATTG GATCCGTTTCG 3251  
 30 CGGGTCCCCT 3261

35 (2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 145 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

45 Met Val Ser Arg Asp Gln Thr Asn Asp Arg Ser Asp Asp Lys Pro Ala  
 1 5 10 15

Arg Ser Asn Pro Thr Asp Cys Ser Val His Thr Glu Pro Ser Asp Ala  
                   20                  25                  30

5 Asn Asn Arg Thr Gly Val His Ser Gly Arg His Pro Gly Glu Ala His  
                   35                  40                  45

Ser Gln Val Arg Asp Leu Asp Leu Gln Phe Asp Cys Gly Gly His Arg  
           50                  55                  60

10 Val Arg Ala Asn Cys Leu Phe Pro Trp Ile Pro Trp Leu Asn Cys Gly  
       65                  70                  75                  80

Cys Ser Leu His Thr Ala Gly Gln Trp Glu Leu Gln Val Arg Ser Asp  
                   85                  90                  95

15 Ala Pro Asp Cys Pro Glu Pro Thr Gly Gln Leu Gln Leu Leu Gln Ala  
                   100                  105                  110

20 Ser Glu Ser Glu Ser His Ser Glu Val Lys His Thr Ser Trp Trp Arg  
           115                  120                  125

Leu Cys Thr Lys Arg His His Lys Arg Arg Asp Leu Pro Arg Lys Pro  
       130                  135                  140

25 Glu  
       145

30 (2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 3261 base pairs

(B) TYPE: nucleic acid

35 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

40

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 131..3166

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

GGATACGATC GGTCTGACCC CGGGGGAGTC ACCCGGGGAC AGGCCGTCAA GGCCTTGTTT

	CAGGATGGGA CTCCTCCTTC TACAACGCTA TCATTGATGG TTAGTAGAGA TCAGACAAAC	120
5	GATCGCAGCG ATG ACA AAC CTG CAA GAT CAA ACC CAA CAG ATT GTT CCG Met Thr Asn Leu Gln Asp Gln Thr Gln Gln Ile Val Pro	169
	1 5 10	
10	TTC ATA CGG AGC CTT CTG ATG CCA ACA ACC GGA CCG GCG TCC ATT CCG Phe Ile Arg Ser Leu Leu Met Pro Thr Thr Gly Pro Ala Ser Ile Pro	217
	15 20 25	
15	GAC GAC ACC CTG GAG AAG CAC ACT CTC AGG TCA GAG ACC TCG ACC TAC Asp Asp Thr Leu Glu Lys His Thr Leu Arg Ser Glu Thr Ser Thr Tyr	265
	30 35 40 45	
	AAT TTG ACT GTG GGG GAC ACA GGG TCA GGG CTA ATT GTC TTT TTC CCT Asn Leu Thr Val Gly Asp Thr Gly Ser Gly Leu Ile Val Phe Phe Pro	313
	50 55 60	
20	GGA TTC CCT GGC TCA ATT GTG GGT GCT CAC TAC ACA CTG CAG GGC AAT Gly Phe Pro Gly Ser Ile Val Gly Ala His Tyr Thr Leu Gln Gly Asn	361
	65 70 75	
25	GGG AAC TAC AAG TTC GAT CAG ATG CTC CTG ACT GCC CAG AAC CTA CCG Gly Asn Tyr Lys Phe Asp Gln Met Leu Leu Thr Ala Gln Asn Leu Pro	409
	80 85 90	
30	GCC AGT TAC AAC TAC TGC AGG CTA GTG AGT CCG AGT CTC ACA GTG AGG Ala Ser Tyr Asn Tyr Cys Arg Leu Val Ser Arg Ser Leu Thr Val Arg	457
	95 100 105	
35	TCA AGC ACA CTT CCT GGT GGC GTT TAT GCA CTA AAC GGC ACC ATA AAC Ser Ser Thr Leu Pro Gly Gly Val Tyr Ala Leu Asn Gly Thr Ile Asn	505
	110 115 120 125	
	GCC GTG ACC TTC CAA GGA AGC CTG AGT GAA CTG ACA GAT GTT AGC TAC Ala Val Thr Phe Gln Gly Ser Leu Ser Glu Leu Thr Asp Val Ser Tyr	553
	130 135 140	
40	AAT GGG TTG ATG TCT GCA ACA GCC AAC ATC AAC GAC AAA ATT GGG AAC Asn Gly Leu Met Ser Ala Thr Ala Asn Ile Asn Asp Lys Ile Gly Asn	601
	145 150 155	
45	GTC CTA GTA GGG GAA GGG GTC ACC GTC CTC AGC TTA CCC ACA TCA TAT Val Leu Val Gly Glu Gly Val Thr Val Leu Ser Leu Pro Thr Ser Tyr	649
	160 165 170	
	GAT CTT GGG TAT GTG AGG CTT GGT GAC CCC ATT CCC GCA ATA GGG CTT	697

Asp Leu Gly Tyr Val Arg Leu Gly Asp Pro Ile Pro Ala Ile Gly Leu  
175 180 185

5 GAC CCA AAA ATG GTA GCC ACA TGT GAC AGC AGT GAC AGG CCC AGA GTC 745  
Asp Pro Lys Met Val Ala Thr Cys Asp Ser Ser Asp Arg Pro Arg Val  
190 195 200 205

10 TAC ACC ATA ACT GCA GCC GAT GAT TAC CAA TTC TCA TCA CAG TAC CAA 793  
Tyr Thr Ile Thr Ala Ala Asp Asp Tyr Gln Phe Ser Ser Gln Tyr Gln  
210 215 220

CCA GGT GGG GTA ACA ATC ACA CTG TTC TCA GCC AAC ATT GAT GCC ATC  
Pro Gly Gly Val Thr Ile Thr Leu Phe Ser Ala Asn Ile Asp Ala Ile 841  
225 230 235

ACA AGC CTC AGC GTT GGG GGA GAG CTC GTG TTT CAA ACA AGC GTC CAC  
Thr Ser Leu Ser Val Gly Gly Glu Leu Val Phe Gln Thr Ser Val His 889  
240 245 250

20 GGC CTT GTA CTG GGC GCC ACC ATC TAC CTC ATA GGC TTT GAT GGG ACA  
Gly Leu Val Leu Gly Ala Thr Ile Tyr Leu Ile Gly Phe Asp Gly Thr 937  
255 260 265

25	ACG GTA ATC ACC AGG GCT GTG GCC GCA AAC AAT GGG CTG ACG ACC GGC	
	Thr Val Ile Thr Arg Ala Val Ala Ala Asn Asn Gly Leu Thr Thr Gly	985
	270                          275                          280                          285	

ACC GAC AAC CTT ATG CCA TTC AAT CTT GTG ATT CCA ACA AAC GAG ATA 1033  
Thr Asp Asn Leu Met Pro Phe Asn Leu Val Ile Pro Thr Asn Glu Ile  
290 295 300

ACC CAG CCA ATC ACA TCC ATC AAA CTG GAG ATA GTG ACC TCC AAA AGT  
Thr Gln Pro Ile Thr Ser Ile Lys Leu Glu Ile Val Thr Ser Lys Ser 1081  
305 310 315

GGT GGT CAG GCA GGG GAT CAG ATG TCA TGG TCG GCA AGA GGG AGC CTA  
Gly Gly Gln Ala Gly Asp Gln Met Ser Trp Ser Ala Arg Gly Ser Leu 1129  
320 325 330

40 GCA GTG ACG ATC CAT GGT GGC AAC TAT CCA GGG GCC CTC CGT CCC GTC 1177  
Ala Val Thr Ile His Gly Gly Asn Tyr Pro Gly Ala Leu Arg Pro Val  
335 340 345

45	ACG	CTA	GTG	GCC	TAC	GAA	AGA	GTG	GCA	ACA	GGA	TCC	GTC	GTT	ACG	GTC	1225
	Thr	Leu	Val	Ala	Tyr	Glu	Arg	Val	Ala	Thr	Gly	Ser	Val	Val	Thr	Val	
	350					355					360					365	

GCT GGG GTG AGC AAC TTC GAG CTG ATC CCA AAT CCT GAA CTA GCA AAG 1273

	Ala Gly Val Ser Asn Phe Glu Leu Ile Pro Asn Pro Glu Leu Ala Lys	
	370 375 380	
5	AAC CTG GTT ACA GAA TAC GGC CGA TTT GAC CCA GGA GCC ATG AAC TAC Asn Leu Val Thr Glu Tyr Gly Arg Phe Asp Pro Gly Ala Met Asn Tyr	1321
	385 390 395	
10	ACA AAA TTG ATA CTG AGT GAG AGG GAC CGT CTT GGC ATC AAG ACC GTC Thr Lys Leu Ile Leu Ser Glu Arg Asp Arg Leu Gly Ile Lys Thr Val	1369
	400 405 410	
15	TGG CCA ACA AGG GAG TAC ACT GAC TTT CGT GAA TAC TTC ATG GAG GTG Trp Pro Thr Arg Glu Tyr Thr Asp Phe Arg Glu Tyr Phe Met Glu Val	1417
	415 420 425	
20	GCC GAC CTC AAC TCT CCC CTG AAG ATT GCA GGA GCA TTC GGC TTC AAA Ala Asp Leu Asn Ser Pro Leu Lys Ile Ala Gly Ala Phe Gly Phe Lys	1465
	430 435 440 445	
25	GAC ATA ATC CGG GCC ATA AGG AGG ATA GCT GTG CCG GTG GTC TCC ACA Asp Ile Ile Arg Ala Ile Arg Arg Ile Ala Val Pro Val Val Ser Thr	1513
	450 455 460	
30	TTG TTC CCA CCT GCC GCT CCC CTA GCC CAT GCA ATT GGG GAA GGT GTA Leu Phe Pro Pro Ala Ala Pro Leu Ala His Ala Ile Gly Glu Gly Val	1561
	465 470 475	
35	GAC TAC CTG CTG GGC GAT GAG GCA CAG GCT GCT TCA GGA ACT GCT CGA Asp Tyr Leu Leu Gly Asp Glu Ala Gln Ala Ala Ser Gly Thr Ala Arg	1609
	480 485 490	
40	GCC GCG TCA GGA AAA GCA AGA GCT GCC TCA GGC CGC ATA AGG CAG CTG Ala Ala Ser Gly Lys Ala Arg Ala Ala Ser Gly Arg Ile Arg Gln Leu	1657
	495 500 505	
45	ACT CTC GCC GCC GAC AAG GGG TAC GAG GTA GTC GCG AAT CTA TTC CAG Thr Leu Ala Ala Asp Lys Gly Tyr Glu Val Val Ala Asn Leu Phe Gln	1705
	510 515 520 525	
50	GTG CCC CAG AAT CCC GTA GTC GAC GGG ATT CTT GCT TCA CCT GGG GTA Val Pro Gln Asn Pro Val Val Asp Gly Ile Leu Ala Ser Pro Gly Val	1753
	530 535 540	
55	CTC CGC GGT GCA CAC AAC CTC GAC TGC GTG TTA AGA GAG GGT GCC ACG Leu Arg Gly Ala His Asn Leu Asp Cys Val Leu Arg Glu Gly Ala Thr	1801
	545 550 555	
60	CTA TTC CCT GTG GTT ATT ACG ACA GTG GAA GAC GCC ATG ACA CCC AAA	1849

	Leu	Phe	Pro	Val	Val	Ile	Thr	Thr	Val	Glu	Asp	Ala	Met	Thr	Pro	Lys	
			560						565					570			
	GCA	TTG	AAC	AGC	AAA	ATG	TTT	GCT	GTC	ATT	GAA	GGC	GTG	CGA	GAA	GAC	1897
5	Ala	Leu	Asn	Ser	Lys	Met	Phe	Ala	Val	Ile	Glu	Gly	Val	Arg	Glu	Asp	
		575					580					585					
	CTC	CAA	CCT	CCA	TCT	CAA	AGA	GGA	TCC	TTC	ATA	CGA	ACT	CTC	TCT	GGA	1945
	Leu	Gln	Pro	Pro	Ser	Gln	Arg	Gly	Ser	Phe	Ile	Arg	Thr	Leu	Ser	Gly	
10	590					595					600					605	
	CAC	AGA	GTC	TAT	GGA	TAT	GCT	CCA	GAT	GGG	GTA	CTT	CCA	CTG	GAG	ACT	1993
	His	Arg	Val	Tyr	Gly	Tyr	Ala	Pro	Asp	Gly	Val	Leu	Pro	Leu	Glu	Thr	
					610					615					620		
15	GGG	AGA	GAC	TAC	ACC	GTT	GTC	CCA	ATA	GAT	GAT	GTC	TGG	GAC	GAC	AGC	2041
	Gly	Arg	Asp	Tyr	Thr	Val	Val	Pro	Ile	Asp	Asp	Val	Trp	Asp	Asp	Ser	
				625					630					635			
	ATT	ATG	CTG	TCC	AAA	GAT	CCC	ATA	CCT	CCT	ATT	GTG	GGA	AAC	AGT	GGA	2089
20	Ile	Met	Leu	Ser	Lys	Asp	Pro	Ile	Pro	Pro	Ile	Val	Gly	Asn	Ser	Gly	
			640					645					650				
	AAT	CTA	GCC	ATA	GCT	TAC	ATG	GAT	GTG	TTT	CGA	CCC	AAA	GTC	CCA	ATC	2137
25	Asn	Leu	Ala	Ile	Ala	Tyr	Met	Asp	Val	Phe	Arg	Pro	Lys	Val	Pro	Ile	
		655					660					665					
	CAT	GTG	GCT	ATG	ACG	GGA	GCC	CTC	AAT	GCT	TGT	GGC	GAG	ATT	GAG	AAA	2185
	His	Val	Ala	Met	Thr	Gly	Ala	Leu	Asn	Ala	Cys	Gly	Glu	Ile	Glu	Lys	
30	670					675					680					685	
	GTA	AGC	TTT	AGA	AGC	ACC	AAG	CTC	GCC	ACT	GCA	CAC	CGA	CTT	GGC	CTT	2233
	Val	Ser	Phe	Arg	Ser	Thr	Lys	Leu	Ala	Thr	Ala	His	Arg	Leu	Gly	Leu	
					690					695					700		
35	AGG	TTG	GCT	GGT	CCC	GGA	GCA	TTC	GAT	GTA	AAC	ACC	GGG	CCC	AAC	TGG	2281
	Arg	Leu	Ala	Gly	Pro	Gly	Ala	Phe	Asp	Val	Asn	Thr	Gly	Pro	Asn	Trp	
				705					710					715			
	GCA	ACG	TTC	ATC	AAA	CGT	TTC	CCT	CAC	AAT	CCA	CGC	GAC	TGG	GAC	AGG	2329
40	Ala	Thr	Phe	Ile	Lys	Arg	Phe	Pro	His	Asn	Pro	Arg	Asp	Trp	Asp	Arg	
			720					725					730				
	CTC	CCC	TAC	CTC	AAC	CTA	CCA	TAC	CTT	CCA	CCC	AAT	GCA	GGA	CGC	CAG	2377
45	Leu	Pro	Tyr	Leu	Asn	Leu	Pro	Tyr	Leu	Pro	Pro	Asn	Ala	Gly	Arg	Gln	
		735					740					745					
	TAC	CAC	CTT	GCC	ATG	GCT	GCA	TCA	GAG	TTC	AAA	GAG	ACC	CCC	GAA	CTC	2425

Tyr His Leu Ala Met Ala Ala Ser Glu Phe Lys Glu Thr Pro Glu Leu  
750 755 760 765

5 GAG AGT GCC GTC AGA GCA ATG GAA GCA GCA GCC AAC GTG GAC CCA CTA 2473  
Glu Ser Ala Val Arg Ala Met Glu Ala Ala Ala Asn Val Asp Pro Leu  
770 775 780

10      TTC CAA TCT GCA CTC AGT GTG TTC ATG TGG CTG GAA GAG AAT GGG ATT      2521  
Phe Gln Ser Ala Leu Ser Val Phe Met Trp Leu Glu Glu Asn Gly Ile  
              785                     790                             795

GTG ACT GAC ATG GCC AAC TTC GCA CTC AGC GAC CCG AAC GCC CAT CGG 2569  
Val Thr Asp Met Ala Asn Phe Ala Leu Ser Asp Pro Asn Ala His Arg  
800 805 810

15  
ATG CGA AAT TTT CTT GCA AAC GCA CCA CAA GCA GGC AGC AAG TCG CAA  
Met Arg Asn Phe Leu Ala Asn Ala Pro Gln Ala Gly Ser Lys Ser Gln  
815 820 825 2617

20 AGG GCC AAG TAC GGG ACA GCA GGC TAC GGA GTG GAG GCT CGG GGC CCC 2665  
Arg Ala Lys Tyr Gly Thr Ala Gly Tyr Gly Val Glu Ala Arg Gly Pro  
830 835 840 845

[illegible]

30 ATG GAG ACC ATG GGC ATC TAC TTT GCA ACA CCA GAA TGG GTA GCA CTC 2761  
Met Glu Thr Met Gly Ile Tyr Phe Ala Thr Pro Glu Trp Val Ala Leu  
865 870 875

AAT GGG CAC CGA GGG CCA AGC CCC GGC CAG CTA AAG TAC TGG CAG AAC 2809  
 Asn Gly His Arg Gly Pro Ser Pro Gly Gln Leu Lys Tyr Trp Gln Asn  
 880 885 890

35

ACA CGA GAA ATA CCG GAC CCA AAC GAG GAC TAT CTA GAC TAC GTG CAT 2857  
Thr Arg Glu Ile Pro Asp Pro Asn Glu Asp Tyr Leu Asp Tyr Val His  
895 900 905

40 GCA GAG AAG AGC CGG TTG GCA TCA GAA GAA CAA ATC CTA AGG GCA GCT 2905  
Ala Glu Lys Ser Arg Leu Ala Ser Glu Glu Gln Ile Leu Arg Ala Ala  
910 915 920 925

45    ACG TCG ATC TAC GGG GCT CCA GGA CAG GCA GAG CCA CCC CAA GCT TTC                  2953  
     Thr Ser Ile Tyr Gly Ala Pro Gly Gln Ala Glu Pro Pro Gln Ala Phe  
                    930                                  935    940

ATA GAC GAA GTT GCC AAA GTC TAT GAA ATC AAC CAT GGA CGT GGC CCA 3001

Ile Asp Glu Val Ala Lys Val Tyr Glu Ile Asn His Gly Arg Gly Pro  
945 950 955

5 AAC CAA GAA CAG ATG AAA GAT CTG CTC TTG ACT GCG ATG GAG ATG AAG 3049  
Asn Gln Glu Gln Met Lys Asp Leu Leu Leu Thr Ala Met Glu Met Lys  
960 965 970

10 CAT CGC AAT CCC AGG CGG GCT CTA CCA AAG CCC AAG CCA AAA CCC AAT 3097  
His Arg Asn Pro Arg Arg Ala Leu Pro Lys Pro Lys Pro Lys Pro Asn  
975 980 985

GCT CCA ACA CAG AGA CCC CCT GGT CGG CTG GGC CGC TGG ATC AGG ACC 3145  
Ala Pro Thr Gln Arg Pro Pro Gly Arg Leu Gly Arg Trp Ile Arg Thr  
990 995 1000 1005

GTC TCT GAT GAG GAC CTT GAG TGAGGCTCCT GGGAGTCTCC CGACACCACC 3196  
Val Ser Asp Glu Asp Leu Glu  
1010

20	CGCGCAGGTG TGGACACCAA TTCGGCCTTA CAACATCCCA AATTGGATCC GTTCGCGGGT	3256
	CCCCT	3261

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1012 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

Met Thr Asn Leu Gln Asp Gln Thr Gln Gln Ile Val Pro Phe Ile Arg  
1 5 10 15

40    Ser Leu Leu Met Pro Thr Thr Gly Pro Ala Ser Ile Pro Asp Asp Thr  
                20                    25                    30

Leu Glu Lys His Thr Leu Arg Ser Glu Thr Ser Thr Tyr Asn Leu Thr  
35 40 45

45 Val Gly Asp Thr Gly Ser Gly Leu Ile Val Phe Phe Pro Gly Phe Pro  
50 55 60

Gly Ser Ile Val Gly Ala His Tyr Thr Leu Gln Gly Asn Gly Asn Tyr



65

70

75

80

Lys Phe Asp Gln Met Leu Leu Thr Ala Gln Asn Leu Pro Ala Ser Tyr  
                             85                            90                            95

5

Asn Tyr Cys Arg Leu Val Ser Arg Ser Leu Thr Val Arg Ser Ser Thr  
                             100                            105                            110

10

Leu Pro Gly Gly Val Tyr Ala Leu Asn Gly Thr Ile Asn Ala Val Thr  
                             115                            120                            125

Phe Gln Gly Ser Leu Ser Glu Leu Thr Asp Val Ser Tyr Asn Gly Leu  
                             130                            135                            140

15

Met Ser Ala Thr Ala Asn Ile Asn Asp Lys Ile Gly Asn Val Leu Val  
                             145                            150                            155                            160

Gly Glu Gly Val Thr Val Leu Ser Leu Pro Thr Ser Tyr Asp Leu Gly  
                             165                            170                            175

20

Tyr Val Arg Leu Gly Asp Pro Ile Pro Ala Ile Gly Leu Asp Pro Lys  
                             180                            185                            190

25

Met Val Ala Thr Cys Asp Ser Ser Asp Arg Pro Arg Val Tyr Thr Ile  
                             195                            200                            205

Thr Ala Ala Asp Asp Tyr Gln Phe Ser Ser Gln Tyr Gln Pro Gly Gly  
                             210                            215                            220

30

Val Thr Ile Thr Leu Phe Ser Ala Asn Ile Asp Ala Ile Thr Ser Leu  
                             225                            230                            235                            240

Ser Val Gly Gly Glu Leu Val Phe Gln Thr Ser Val His Gly Leu Val  
                             245                            250                            255

35

Leu Gly Ala Thr Ile Tyr Leu Ile Gly Phe Asp Gly Thr Thr Val Ile  
                             260                            265                            270

40

Thr Arg Ala Val Ala Ala Asn Asn Gly Leu Thr Thr Gly Thr Asp Asn  
                             275                            280                            285

Leu Met Pro Phe Asn Leu Val Ile Pro Thr Asn Glu Ile Thr Gln Pro  
                             290                            295                            300

45

Ile Thr Ser Ile Lys Leu Glu Ile Val Thr Ser Lys Ser Gly Gly Gln  
                             305                            310                            315                            320

Ala Gly Asp Gln Met Ser Trp Ser Ala Arg Gly Ser Leu Ala Val Thr

	325	330	335
	Ile His Gly Gly Asn Tyr Pro Gly Ala Leu Arg Pro Val Thr Leu Val		
	340	345	350
5	Ala Tyr Glu Arg Val Ala Thr Gly Ser Val Val Thr Val Ala Gly Val		
	355	360	365
	Ser Asn Phe Glu Leu Ile Pro Asn Pro Glu Leu Ala Lys Asn Leu Val		
10	370	375	380
	Thr Glu Tyr Gly Arg Phe Asp Pro Gly Ala Met Asn Tyr Thr Lys Leu		
	385	390	395 400
15	Ile Leu Ser Glu Arg Asp Arg Leu Gly Ile Lys Thr Val Trp Pro Thr		
	405	410	415
	Arg Glu Tyr Thr Asp Phe Arg Glu Tyr Phe Met Glu Val Ala Asp Leu		
	420	425	430
20	Asn Ser Pro Leu Lys Ile Ala Gly Ala Phe Gly Phe Lys Asp Ile Ile		
	435	440	445
	Arg Ala Ile Arg Arg Ile Ala Val Pro Val Val Ser Thr Leu Phe Pro		
25	450	455	460
	Pro Ala Ala Pro Leu Ala His Ala Ile Gly Glu Gly Val Asp Tyr Leu		
	465	470	475 480
30	Leu Gly Asp Glu Ala Gln Ala Ala Ser Gly Thr Ala Arg Ala Ala Ser		
	485	490	495
	Gly Lys Ala Arg Ala Ala Ser Gly Arg Ile Arg Gln Leu Thr Leu Ala		
	500	505	510
35	Ala Asp Lys Gly Tyr Glu Val Val Ala Asn Leu Phe Gln Val Pro Gln		
	515	520	525
	Asn Pro Val Val Asp Gly Ile Leu Ala Ser Pro Gly Val Leu Arg Gly		
40	530	535	540
	Ala His Asn Leu Asp Cys Val Leu Arg Glu Gly Ala Thr Leu Phe Pro		
	545	550	555 560
45	Val Val Ile Thr Thr Val Glu Asp Ala Met Thr Pro Lys Ala Leu Asn		
	565	570	575
	Ser Lys Met Phe Ala Val Ile Glu Gly Val Arg Glu Asp Leu Gln Pro		

	580	585	590
	Pro Ser Gln Arg Gly Ser Phe Ile Arg Thr Leu Ser Gly His Arg Val		
	595	600	605
5	Tyr Gly Tyr Ala Pro Asp Gly Val Leu Pro Leu Glu Thr Gly Arg Asp		
	610	615	620
10	Tyr Thr Val Val Pro Ile Asp Asp Val Trp Asp Asp Ser Ile Met Leu		
	625	630	635
	Ser Lys Asp Pro Ile Pro Pro Ile Val Gly Asn Ser Gly Asn Leu Ala		
	645	650	655
15	Ile Ala Tyr Met Asp Val Phe Arg Pro Lys Val Pro Ile His Val Ala		
	660	665	670
	Met Thr Gly Ala Leu Asn Ala Cys Gly Glu Ile Glu Lys Val Ser Phe		
	675	680	685
20	Arg Ser Thr Lys Leu Ala Thr Ala His Arg Leu Gly Leu Arg Leu Ala		
	690	695	700
	Gly Pro Gly Ala Phe Asp Val Asn Thr Gly Pro Asn Trp Ala Thr Phe		
25	705	710	715
	Ile Lys Arg Phe Pro His Asn Pro Arg Asp Trp Asp Arg Leu Pro Tyr		
	725	730	735
30	Leu Asn Leu Pro Tyr Leu Pro Pro Asn Ala Gly Arg Gln Tyr His Leu		
	740	745	750
	Ala Met Ala Ala Ser Glu Phe Lys Glu Thr Pro Glu Leu Glu Ser Ala		
	755	760	765
35	Val Arg Ala Met Glu Ala Ala Ala Asn Val Asp Pro Leu Phe Gln Ser		
	770	775	780
	Ala Leu Ser Val Phe Met Trp Leu Glu Glu Asn Gly Ile Val Thr Asp		
40	785	790	795
	Met Ala Asn Phe Ala Leu Ser Asp Pro Asn Ala His Arg Met Arg Asn		
	805	810	815
45	Phe Leu Ala Asn Ala Pro Gln Ala Gly Ser Lys Ser Gln Arg Ala Lys		
	820	825	830
	Tyr Gly Thr Ala Gly Tyr Gly Val Glu Ala Arg Gly Pro Thr Pro Glu		

	835	840	845
	Glu Ala Gln Arg Glu Lys Asp Thr Arg Ile Ser Lys Lys Met Glu Thr		
	850	855	860
5	Met Gly Ile Tyr Phe Ala Thr Pro Glu Trp Val Ala Leu Asn Gly His		
	865	870	875 880
	Arg Gly Pro Ser Pro Gly Gln Leu Lys Tyr Trp Gln Asn Thr Arg Glu		
10		885	890 895
	Ile Pro Asp Pro Asn Glu Asp Tyr Leu Asp Tyr Val His Ala Glu Lys		
		900	905 910
15	Ser Arg Leu Ala Ser Glu Glu Gln Ile Leu Arg Ala Ala Thr Ser Ile		
		915	920 925
	Tyr Gly Ala Pro Gly Gln Ala Glu Pro Pro Gln Ala Phe Ile Asp Glu		
		930	935 940
20	Val Ala Lys Val Tyr Glu Ile Asn His Gly Arg Gly Pro Asn Gln Glu		
		945	950 955 960
	Gln Met Lys Asp Leu Leu Leu Thr Ala Met Glu Met Lys His Arg Asn		
25		965	970 975
	Pro Arg Arg Ala Leu Pro Lys Pro Lys Pro Lys Pro Asn Ala Pro Thr		
		980	985 990
30	Gln Arg Pro Pro Gly Arg Leu Gly Arg Trp Ile Arg Thr Val Ser Asp		
		995	1000 1005
	Glu Asp Leu Glu		
	1010		

35

## (2) INFORMATION FOR SEQ ID NO: 7:

40

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 3261 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

45

## (ii) MOLECULE TYPE: cDNA

## (ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION:97..531

5

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

GGATACGATC GGTCTGACCC CGGGGGAGTC ACCCGGGGAC AGGCCGTCAA GGCCTTGTTTC 60

10 CAGGATGGGA CTCCTCCTTC TACAACGCTA TCATTG GAA GTT AGT TGA GAT CTG 114  
 Glu Val Ser \* Asp Leu  
 1 5

15 ACA AAC GAT CGC AGC GAT GAC AAA CCT GCA AGA TCA AAC CCA ACA GAT 162  
 Thr Asn Asp Arg Ser Asp Asp Lys Pro Ala Arg Ser Asn Pro Thr Asp  
 10 15 20

## 20 (2) INFORMATION FOR SEQ ID NO: 8:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 2827 base pairs

(B) TYPE: nucleic acid

25 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

30

## (ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION:112..2745

35

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

GGATACGATG GGTCTGACCC TCTGGGAGTC ACGAATTAAC GTGGCTACTA GGGGCGATAC 60

40 CCGCCGCTGG CTGCCACGTT AGTGGCTCCT CTTCTTGATG ATTCTGCCAC C ATG AGT 117  
 Met Ser  
 1

45 GAC ATT TTC AAC AGT CCA CAG GCG CGA AGC ACG ATC TCA GCA GCG TTC 165  
 Asp Ile Phe Asn Ser Pro Gln Ala Arg Ser Thr Ile Ser Ala Ala Phe  
 5 10 15

GGC ATA AAG CCT ACT GCT GGA CAA GAC GTG GAA GAA CTC TTG ATC CCT 213

Gly Ile Lys Pro Thr Ala Gly Gln Asp Val Glu Glu Leu Leu Ile Pro  
 20 25 30

5 AAA GTT TGG GTG CCA CCT GAG GAT CCG CTT GCC AGC CCT AGT CGA CTG 261  
 Lys Val Trp Val Pro Pro Glu Asp Pro Leu Ala Ser Pro Ser Arg Leu  
 35 40 45 50

10 GCA AAG TTC CTC AGA GAG AAC GGC TAC AAA GTT TTG CAG CCG CGG TCT 309  
 Ala Lys Phe Leu Arg Glu Asn Gly Tyr Lys Val Leu Gln Pro Arg Ser  
 55 60 65

15 CTG CCC GAG AAT GAG GAG TAT GAG ACC GAC CAA ATA CTC CCA GAC TTA 357  
 Leu Pro Glu Asn Glu Glu Tyr Glu Thr Asp Gln Ile Leu Pro Asp Leu  
 70 75 80

GCA TGG ATG CGA CAG ATA GAA GGG GCT GTT TTA AAA CCC ACT CTA TCT 405  
 Ala Trp Met Arg Gln Ile Glu Gly Ala Val Leu Lys Pro Thr Leu Ser  
 85 90 95

20 CTC CCT ATT GGA GAT CAG GAG TAC TTC CCA AAG TAC TAC CCA ACA CAT 453  
 Leu Pro Ile Gly Asp Gln Glu Tyr Phe Pro Lys Tyr Tyr Pro Thr His  
 100 105 110

25 CGC CCT AGC AAG GAG AAG CCC AAT GCG TAC CCG CCA GAC ATC GCA CTA 501  
 Arg Pro Ser Lys Glu Lys Pro Asn Ala Tyr Pro Pro Asp Ile Ala Leu  
 115 120 125 130

30 CTC AAG CAG ATG ATT TAC CTG TTT CTC CAG GTT CCA GAG GCC AAC GAG 549  
 Leu Lys Gln Met Ile Tyr Leu Phe Leu Gln Val Pro Glu Ala Asn Glu  
 135 140 145

35 GGC CTA AAG GAT GAA GTA ACC CTC TTG ACC CAA AAC ATA AGG GAC AAG 597  
 Gly Leu Lys Asp Glu Val Thr Leu Leu Thr Gln Asn Ile Arg Asp Lys  
 150 155 160

GCC TAT GGA AGT GGG ACC TAC ATG GGA CAA GCA ACT CGA CTT GTG GCC 645  
 Ala Tyr Gly Ser Gly Thr Tyr Met Gly Gln Ala Thr Arg Leu Val Ala  
 165 170 175

40 ATG AAG GAG GTC GCC ACT GGA AGA AAC CCA AAC AAG GAT CCT CTA AAG 693  
 Met Lys Glu Val Ala Thr Gly Arg Asn Pro Asn Lys Asp Pro Leu Lys  
 180 185 190

45 CTT GGG TAC ACT TTT GAG AGC ATC GCG CAG CTA CTT GAC ATC ACA CTA 741  
 Leu Gly Tyr Thr Phe Glu Ser Ile Ala Gln Leu Leu Asp Ile Thr Leu  
 195 200 205 210

CCG GTA GGC CCA CCC GGT GAG GAT GAC AAG CCC TGG GTG CCA CTC ACA 789

	Pro	Val	Gly	Pro	Pro	Gly	Glu	Asp	Asp	Lys	Pro	Trp	Val	Pro	Leu	Thr	
					215					220					225		
	AGA	GTG	CCG	TCA	CGG	ATG	TTG	GTG	CTG	ACG	GGA	GAC	GTA	GAT	GGC	GAC	837
5	Arg	Val	Pro	Ser	Arg	Met	Leu	Val	Leu	Thr	Gly	Asp	Val	Asp	Gly	Asp	
				230					235					240			
	TTT	GAG	GTT	GAA	GAT	TAC	CTT	CCC	AAA	ATC	AAC	CTC	AAG	TCA	TCA	AGT	885
10	Phe	Glu	Val	Glu	Asp	Tyr	Leu	Pro	Lys	Ile	Asn	Leu	Lys	Ser	Ser	Ser	
			245					250					255				
	GGA	CTA	CCA	TAT	GTA	GGT	CGC	ACC	AAA	GGA	GAG	ACA	ATT	GGC	GAG	ATG	933
15	Gly	Leu	Pro	Tyr	Val	Gly	Arg	Thr	Lys	Gly	Glu	Thr	Ile	Gly	Glu	Met	
		260					265					270					
	ATA	GCT	ATA	TCA	AAC	CAG	TTT	CTC	AGA	GAG	CTA	TCA	ACA	CTG	TTG	AAG	981
	Ile	Ala	Ile	Ser	Asn	Gln	Phe	Leu	Arg	Glu	Leu	Ser	Thr	Leu	Leu	Lys	
	275					280				285						290	
20	CAA	GGT	GCA	GGG	ACA	AAG	GGG	TCA	AAC	AAG	AAG	AAG	CTA	CTC	AGC	ATG	1029
	Gln	Gly	Ala	Gly	Thr	Lys	Gly	Ser	Asn	Lys	Lys	Lys	Leu	Leu	Ser	Met	
					295					300					305		
	TTA	AGT	GAC	TAT	TGG	TAC	TTA	TCA	TGC	GGG	CTT	TTG	TTT	CCA	AAG	GCT	1077
25	Leu	Ser	Asp	Tyr	Trp	Tyr	Leu	Ser	Cys	Gly	Leu	Leu	Phe	Pro	Lys	Ala	
				310					315					320			
	GAA	AGG	TAC	GAC	AAA	AGT	ACA	TGG	CTC	ACC	AAG	ACC	CGG	AAC	ATA	TGG	1125
30	Glu	Arg	Tyr	Asp	Lys	Ser	Thr	Trp	Leu	Thr	Lys	Thr	Arg	Asn	Ile	Trp	
			325					330					335				
	TCA	GCT	CCA	TCC	CCA	ACA	CAC	CTC	ATG	ATC	TCC	ATG	ATC	ACC	TGG	CCC	1173
35	Ser	Ala	Pro	Ser	Pro	Thr	His	Leu	Met	Ile	Ser	Met	Ile	Thr	Trp	Pro	
		340					345					350					
	GTG	ATG	TCC	AAC	AGC	CCA	AAT	AAC	GTG	TTG	AAC	ATT	GAA	GGG	TGT	CCA	1221
	Val	Met	Ser	Asn	Ser	Pro	Asn	Asn	Val	Leu	Asn	Ile	Glu	Gly	Cys	Pro	
	355					360				365					370		
40	TCA	CTC	TAC	AAA	TTC	AAC	CCG	TTC	AGA	GGA	GGG	TTG	AAC	AGG	ATC	GTC	1269
	Ser	Leu	Tyr	Lys	Phe	Asn	Pro	Phe	Arg	Gly	Gly	Leu	Asn	Arg	Ile	Val	
					375					380					385		
	GAG	TGG	ATA	TTG	GCC	CCG	GAA	GAA	CCC	AAG	GCT	CTT	GTA	TAT	GCG	GAC	1317
45	Glu	Trp	Ile	Leu	Ala	Pro	Glu	Glu	Pro	Lys	Ala	Leu	Val	Tyr	Ala	Asp	
				390					395					400			
	AAC	ATA	TAC	ATT	GTC	CAC	TCA	AAC	ACG	TGG	TAC	TCA	ATT	GAC	CTA	GAG	1365

5	AAG GGT GAG GCA AAC TGC ACT CGC CAA CAC ATG CAA GCC GCA ATG TAC Lys Gly Glu Ala Asn Cys Thr Arg Gln His Met Gln Ala Ala Met Tyr 420 425 430	1413
10	TAC ATA CTC ACC AGA GGG TGG TCA GAC AAC GGC GAC CCA ATG TTC AAT Tyr Ile Leu Thr Arg Gly Trp Ser Asp Asn Gly Asp Pro Met Phe Asn 435 440 445 450	1461
15	CAA ACA TGG GCC ACC TTT GCC ATG AAC ATT GCC CCT GCT CTA GTG GTG Gln Thr Trp Ala Thr Phe Ala Met Asn Ile Ala Pro Ala Leu Val Val 455 460 465	1509
20	GAC TCA TCG TGC CTG ATA ATG AAC CTG CAA ATT AAG ACC TAT GGT CAA Asp Ser Ser Cys Leu Ile Met Asn Leu Gln Ile Lys Thr Tyr Gly Gln 470 475 480	1557
25	GGC AGC GGG AAT GCA GCC ACG TTC ATC AAC AAC CAC CTC TTG AGC ACG Gly Ser Gly Asn Ala Ala Thr Phe Ile Asn Asn His Leu Leu Ser Thr 485 490 495	1605
30	CTA GTG CTT GAC CAG TGG AAC TTG ATG AGA CAG CCC AGA CCA GAC AGC Leu Val Leu Asp Gln Trp Asn Leu Met Arg Gln Pro Arg Pro Asp Ser 500 505 510	1653
35	GAG GAG TTC AAA TCA ATT GAG GAC AAG CTA GGT ATC AAC TTT AAG ATT Glu Glu Phe Lys Ser Ile Glu Asp Lys Leu Gly Ile Asn Phe Lys Ile 515 520 525 530	1701
40	GAG AGG TCC ATT GAT GAT ATC AGG GGC AAG CTG AGA CAG CTT GTC CTC Glu Arg Ser Ile Asp Asp Ile Arg Gly Lys Leu Arg Gln Leu Val Leu 535 540 545	1749
45	CTT GCA CAA CCA GGG TAC CTG AGT GGG GGG GTT GAA CCA GAA CAA TCC Leu Ala Gln Pro Gly Tyr Leu Ser Gly Gly Val Glu Pro Glu Gln Ser 550 555 560	1797
50	AGC CCA ACT GTT GAG CTT GAC CTA CTA GGG TGG TCA GCT ACA TAC AGC Ser Pro Thr Val Glu Leu Asp Leu Leu Gly Trp Ser Ala Thr Tyr Ser 565 570 575	1845
55	AAA GAT CTC GGG ATC TAT GTG CCG GTG CTT GAC AAG GAA CGC CTA TTT Lys Asp Leu Gly Ile Tyr Val Pro Val Leu Asp Lys Glu Arg Leu Phe 580 585 590	1893
60	TGT TCT GCT GCG TAT CCC AAG GGA GTA GAG AAC AAG AGT CTC AAG TCC	1941



	Cys	Ser	Ala	Ala	Tyr	Pro	Lys	Gly	Val	Glu	Asn	Lys	Ser	Leu	Lys	Ser	
	595					600					605					610	
5	AAA	GTC	GGG	ATC	GAG	CAG	GCA	TAC	AAG	GTA	GTC	AGG	TAT	GAG	GCG	TTG	1989
	Lys	Val	Gly	Ile	Glu	Gln	Ala	Tyr	Lys	Val	Val	Arg	Tyr	Glu	Ala	Leu	
				615						620					625		
10	AGG	TTG	GTA	GGT	GGT	TGG	AAC	TAC	CCA	CTC	CTG	AAC	AAA	GCC	TGC	AAG	2037
	Arg	Leu	Val	Gly	Gly	Trp	Asn	Tyr	Pro	Leu	Leu	Asn	Lys	Ala	Cys	Lys	
				630					635						640		
15	AAT	AAC	GCA	GGC	GCC	GCT	CGG	CGG	CAT	CTG	GAG	GCC	AAG	GGG	TTC	CCA	2085
	Asn	Asn	Ala	Gly	Ala	Ala	Arg	Arg	His	Leu	Glu	Ala	Lys	Gly	Phe	Pro	
				645					650					655			
20	CTC	GAC	GAG	TTC	CTA	GCC	GAG	TGG	TCT	GAG	CTG	TCA	GAG	TTC	GGT	GAG	2133
	Leu	Asp	Glu	Phe	Leu	Ala	Glu	Trp	Ser	Glu	Leu	Ser	Glu	Phe	Gly	Glu	
				660					665					670			
25	GCC	TTC	GAA	GGC	TTC	AAT	ATC	AAG	CTG	ACC	GTA	ACA	TCT	GAG	AGC	CTA	2181
	Ala	Phe	Glu	Gly	Phe	Asn	Ile	Lys	Leu	Thr	Val	Thr	Ser	Glu	Ser	Leu	
						675					680					690	
30	GCC	GAA	CTG	AAC	AAG	CCA	GTA	CCC	CCC	AAG	CCC	CCA	AAT	GTC	AAC	AGA	2229
	Ala	Glu	Leu	Asn	Lys	Pro	Val	Pro	Pro	Lys	Pro	Pro	Asn	Val	Asn	Arg	
						695					700					705	
35	CCA	GTC	AAC	ACT	GGG	GGA	CTC	AAG	GCA	GTC	AGC	AAC	GCC	CTC	AAG	ACC	2277
	Pro	Val	Asn	Thr	Gly	Gly	Leu	Lys	Ala	Val	Ser	Asn	Ala	Leu	Lys	Thr	
					710					715						720	
40	GGT	CGG	TAC	AGG	AAC	GAA	GCC	GGA	CTG	AGT	GGT	CTC	GTC	CTT	CTA	GCC	2325
	Gly	Arg	Tyr	Arg	Asn	Glu	Ala	Gly	Leu	Ser	Gly	Leu	Val	Leu	Leu	Ala	
				725					730							735	
45	ACA	GCA	AGA	AGC	CGT	CTG	CAA	GAT	GCA	GTT	AAG	GCC	AAG	GCA	GAA	GCC	2373
	Thr	Ala	Arg	Ser	Arg	Leu	Gln	Asp	Ala	Val	Lys	Ala	Lys	Ala	Glu	Ala	
				740					745							750	
50	GAG	AAA	CTC	CAC	AAG	TCC	AAG	CCA	GAC	GAC	CCC	GAT	GCA	GAC	TGG	TTC	2421
	Glu	Lys	Leu	His	Lys	Ser	Lys	Pro	Asp	Asp	Pro	Asp	Ala	Asp	Trp	Phe	
						755					760					770	
55	GAA	AGA	TCA	GAA	ACT	CTG	TCA	GAC	CTT	CTG	GAG	AAA	GCC	GAC	ATC	GCC	2469
	Glu	Arg	Ser	Glu	Thr	Leu	Ser	Asp	Leu	Leu	Glu	Lys	Ala	Asp	Ile	Ala	
					775						780					785	
60	AGC	AAG	GTC	GCC	CAC	TCA	GCA	CTC	GTG	GAA	ACA	AGC	GAC	GCC	CTT	GAA	2517

Ser Lys Val Ala His Ser Ala Leu Val Glu Thr Ser Asp Ala Leu Glu  
 790 795 800

5 GCA GTT CAG TCG ACT TCC GTG TAC ACC CCC AAG TAC CCA GAA GTC AAG 2565  
 Ala Val Gln Ser Thr Ser Val Tyr Thr Pro Lys Tyr Pro Glu Val Lys  
 805 810 815

10 AAC CCA CAG ACC GCC TCC AAC CCC GTT GTT GGG CTC CAC CTG CCC GCC 2613  
 Asn Pro Gln Thr Ala Ser Asn Pro Val Val Gly Leu His Leu Pro Ala  
 820 825 830

15 AAG AGA GCC ACC GGT GTC CAG GCC GCT CTT CTC GGA GCA GGA ACG AGC 2661  
 Lys Arg Ala Thr Gly Val Gln Ala Ala Leu Leu Gly Ala Gly Thr Ser  
 835 840 845 850

AGA CCA ATG GGG ATG GAG GCC CCA ACA CGG TCC AAG AAC GCC GTG AAA 2709  
 Arg Pro Met Gly Met Glu Ala Pro Thr Arg Ser Lys Asn Ala Val Lys  
 855 860 865

20 ATG GCC AAA CGG CGG CAA CGC CAA AAG GAG AGC CGC TAACAGCCAT 2755  
 Met Ala Lys Arg Arg Gln Arg Gln Lys Glu Ser Arg  
 870 875

25 GATGGGAACC ACTCAAGAAG AGGACACTAA TCCCAGACCC CGTATCCCCG GCCTTCGCCT 2815

GCGGGGGCCC CC 2827

CLAIMS

- 1 A birnavirus mutant which is not able to produce a native VP5 protein as a result of a mutation in the VP5 gene of the birnavirus genome.
- 2 A birnavirus mutant according to claim 1, characterised in that the mutation is a substitution.
- 3 A birnavirus mutant according to claim 1, characterised in that the mutation is an insertion of a heterologous nucleic acid sequence.
- 4 A birnavirus mutant according to claim 3, characterised in that the heterologous nucleic acid sequence encodes a polypeptide and the heterologous nucleic acid sequence is under the control of an expression control sequence regulating the expression of the sequence in a cell infected with the virus mutant.
- 5 A birnavirus mutant according to claims 1-4, characterised in that the birnavirus is infectious bursal disease virus (IBDV).
- 6 A birnavirus mutant according to claim 5, characterised in that the mutation is in the genome of a virulent field virus.
- 7 A birnavirus mutant according to claim 5, characterised in that the mutation is in the genome of vaccine strain, preferably in vaccine strain D78.
- 8 A birnavirus mutant according to claims 5-7, characterised in that the mutant has a mutated start codon and three stop codons in the 5'-end of the VP5 gene as shown in SEQ ID No: 7.
- 9 A birnavirus according to claims 5-8, characterised in that the IBDV expresses a chimeric VP2 protein comprising virus neutralising epitopes of different antigenic IBDV types.

- 10 A vaccine against a birnavirus infection in animals, characterised in that it comprises a birnavirus mutant according to claims 1-9 and a pharmaceutically acceptable carrier.
- 5 11 A method for determining birnavirus infection in an animal, characterised in that a sample of the animal is examined for the presence of anti-VP5 antibodies.
- 12 A method according to claim 11, characterised in that the method comprises the steps of:
- 10 (i) incubating a sample suspected of containing anti-birnavirus antibodies, with VP5 antigen,
- (ii) allowing the formation of antibody-antigen complex, and
- (ii) detecting the presence of the antibody-antigen complex.
- 15 13 A diagnostic test kit suitable for carrying out a method according to claims 11-12.
- 14 Use of the lack of the expression of native VP5 protein by a birnavirus mutant as a marker to distinguish vaccinated animals from animals infected with naturally-occurring birnavirus.
- 20


ABSTRACT

The present invention provides a birnavirus mutant which is suited as vaccine candidate in eradication control programmes. The mutant is not able to produce a native VP5 protein, and this  
5 feature can be used as a marker to distinguish between animals vaccinated with the VP5 mutant or infected with a naturally-occurring birnavirus.

# Genomic organization of segment A of strain D78 and segment B of strain P2

Figure 1

## D78 segment A



## P2 segment B




Figure 2




Figure 3






Figure 4





Figure 5



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.